

# $\mathbb{Q}$ -forms and the theory of central simple $G$ -algebras

Jan Jongen

RWTH-Aachen

26.03.2010

## Example 1:

Consider  $G = Q_8$  and choose  $\chi \in \text{Irr}(G)$  with  $\chi(1) = 2$ .

## Example 1:

Consider  $G = Q_8$  and choose  $\chi \in \text{Irr}(G)$  with  $\chi(1) = 2$ .

- A representation which affords  $\chi$  is given by

$$\Delta(a) = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}, \quad \Delta(b) = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

## Example 1:

Consider  $G = Q_8$  and choose  $\chi \in \text{Irr}(G)$  with  $\chi(1) = 2$ .

- A representation which affords  $\chi$  is given by

$$\Delta(a) = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}, \quad \Delta(b) = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

- $\mathbb{Q}[i][x_1, x_2]^G = \langle x_1^2 x_2^2, x_1^4 + x_2^4, x_1 x_2^5 - x_2 x_1^5 \rangle_{\mathbb{Q}[i]\text{-alg}}$

## Example 1:

Consider  $G = Q_8$  and choose  $\chi \in \text{Irr}(G)$  with  $\chi(1) = 2$ .

- A representation which affords  $\chi$  is given by

$$\Delta(a) = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}, \quad \Delta(b) = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

- $\mathbb{Q}[i][x_1, x_2]^G = \langle x_1^2 x_2^2, x_1^4 + x_2^4, x_1 x_2^5 - x_2 x_1^5 \rangle_{\mathbb{Q}[i]\text{-alg}}$

## Observations:

## Example 1:

Consider  $G = Q_8$  and choose  $\chi \in \text{Irr}(G)$  with  $\chi(1) = 2$ .

- A representation which affords  $\chi$  is given by

$$\Delta(a) = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}, \quad \Delta(b) = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

- $\mathbb{Q}[i][x_1, x_2]^G = \langle x_1^2 x_2^2, x_1^4 + x_2^4, x_1 x_2^5 - x_2 x_1^5 \rangle_{\mathbb{Q}[i]\text{-alg}}$

## Observations:

- Ring of polynomial invariants has a generating set of rational polynomials although the representation is not rational

## Example 1:

Consider  $G = Q_8$  and choose  $\chi \in \text{Irr}(G)$  with  $\chi(1) = 2$ .

- A representation which affords  $\chi$  is given by

$$\Delta(a) = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}, \quad \Delta(b) = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

- $\mathbb{Q}[i][x_1, x_2]^G = \langle x_1^2 x_2^2, x_1^4 + x_2^4, x_1 x_2^5 - x_2 x_1^5 \rangle_{\mathbb{Q}[i]\text{-alg}}$

## Observations:

- Ring of polynomial invariants has a generating set of rational polynomials although the representation is not rational
- For all  $g \in G$  applying the GALOIS automorphism  $I \mapsto -I$  to  $\Delta(g)$  is afforded by conjugation with  $\Delta(b) \rightsquigarrow \text{Gal}(\mathbb{Q}[i]/\mathbb{Q})$  acts on  $\Delta(G)$

# Motivation

## Example 2:

Consider  $G = \mathrm{PSL}_2(7)$  and choose  $\chi \in \mathrm{Irr}(G)$  with  $\chi(1) = 3$

# Motivation

## Example 2:

Consider  $G = \mathrm{PSL}_2(7)$  and choose  $\chi \in \mathrm{Irr}(G)$  with  $\chi(1) = 3$

- A representation which affords  $\chi$  is given by

$$\Delta(a) = \begin{pmatrix} \zeta_7 & 0 & 0 \\ 0 & \zeta_7^2 & 0 \\ 0 & 0 & \zeta_7^4 \end{pmatrix}, \quad \Delta(b) = \frac{1}{\sqrt{-7}} \begin{pmatrix} \zeta_7^5 - \zeta_7^2 & \alpha & \zeta_7^3 - \zeta_7^4 \\ \alpha & \zeta_7^3 - \zeta_7^4 & \zeta_7^5 - \zeta_7^2 \\ \zeta_7^3 - \zeta_7^4 & \zeta_7^5 - \zeta_7^2 & \alpha \end{pmatrix}$$

with  $\alpha := -1 - \zeta_7^5 - \zeta_7^4 - \zeta_7^3 - \zeta_7^2 - 2\zeta_7$

# Motivation

## Example 2:

Consider  $G = \mathrm{PSL}_2(7)$  and choose  $\chi \in \mathrm{Irr}(G)$  with  $\chi(1) = 3$

- A representation which affords  $\chi$  is given by

$$\Delta(a) = \begin{pmatrix} \zeta_7 & 0 & 0 \\ 0 & \zeta_7^2 & 0 \\ 0 & 0 & \zeta_7^4 \end{pmatrix}, \quad \Delta(b) = \frac{1}{\sqrt{-7}} \begin{pmatrix} \zeta_7^5 - \zeta_7^2 & \alpha & \zeta_7^3 - \zeta_7^4 \\ \alpha & \zeta_7^3 - \zeta_7^4 & \zeta_7^5 - \zeta_7^2 \\ \zeta_7^3 - \zeta_7^4 & \zeta_7^5 - \zeta_7^2 & \alpha \end{pmatrix}$$

with  $\alpha := -1 - \zeta_7^5 - \zeta_7^4 - \zeta_7^3 - \zeta_7^2 - 2\zeta_7$

## Observations:

# Motivation

## Example 2:

Consider  $G = \mathrm{PSL}_2(7)$  and choose  $\chi \in \mathrm{Irr}(G)$  with  $\chi(1) = 3$

- A representation which affords  $\chi$  is given by

$$\Delta(a) = \begin{pmatrix} \zeta_7 & 0 & 0 \\ 0 & \zeta_7^2 & 0 \\ 0 & 0 & \zeta_7^4 \end{pmatrix}, \quad \Delta(b) = \frac{1}{\sqrt{-7}} \begin{pmatrix} \zeta_7^5 - \zeta_7^2 & \alpha & \zeta_7^3 - \zeta_7^4 \\ \alpha & \zeta_7^3 - \zeta_7^4 & \zeta_7^5 - \zeta_7^2 \\ \zeta_7^3 - \zeta_7^4 & \zeta_7^5 - \zeta_7^2 & \alpha \end{pmatrix}$$

with  $\alpha := -1 - \zeta_7^5 - \zeta_7^4 - \zeta_7^3 - \zeta_7^2 - 2\zeta_7$

## Observations:

- $\mathbb{Q}[\zeta_7][x]^G$  is generated by:  $x_1x_2^3 + x_2x_3^3 + x_3x_1^3, 270x_3^2x_1^2x_2^2 - 54x_3^5x_1 - 54x_2^5x_3 - 54x_1^5x_2, f_3, f_4$  where  $f_3, f_4 \in \mathbb{Q}[x] \rightsquigarrow$   
Generating set of rational invariants!

# Motivation

## Example 2:

Consider  $G = \mathrm{PSL}_2(7)$  and choose  $\chi \in \mathrm{Irr}(G)$  with  $\chi(1) = 3$

- A representation which affords  $\chi$  is given by

$$\Delta(a) = \begin{pmatrix} \zeta_7 & 0 & 0 \\ 0 & \zeta_7^2 & 0 \\ 0 & 0 & \zeta_7^4 \end{pmatrix}, \quad \Delta(b) = \frac{1}{\sqrt{-7}} \begin{pmatrix} \zeta_7^5 - \zeta_7^2 & \alpha & \zeta_7^3 - \zeta_7^4 \\ \alpha & \zeta_7^3 - \zeta_7^4 & \zeta_7^5 - \zeta_7^2 \\ \zeta_7^3 - \zeta_7^4 & \zeta_7^5 - \zeta_7^2 & \alpha \end{pmatrix}$$

with  $\alpha := -1 - \zeta_7^5 - \zeta_7^4 - \zeta_7^3 - \zeta_7^2 - 2\zeta_7$

## Observations:

- $\mathbb{Q}[\zeta_7][x]^G$  is generated by:  $x_1x_2^3 + x_2x_3^3 + x_3x_1^3, 270x_3^2x_1^2x_2^2 - 54x_3^5x_1 - 54x_2^5x_3 - 54x_1^5x_2, f_3, f_4$  where  $f_3, f_4 \in \mathbb{Q}[x] \rightsquigarrow$   
Generating set of rational invariants!
- $\mathrm{Gal}(\mathbb{Q}[\zeta_7]/\mathbb{Q})$  acts on  $\Delta(G)$

# Motivation

## Example 2:

Consider  $G = \mathrm{PSL}_2(7)$  and choose  $\chi \in \mathrm{Irr}(G)$  with  $\chi(1) = 3$

- A representation which affords  $\chi$  is given by

$$\Delta(a) = \begin{pmatrix} \zeta_7 & 0 & 0 \\ 0 & \zeta_7^2 & 0 \\ 0 & 0 & \zeta_7^4 \end{pmatrix}, \quad \Delta(b) = \frac{1}{\sqrt{-7}} \begin{pmatrix} \zeta_7^5 - \zeta_7^2 & \alpha & \zeta_7^3 - \zeta_7^4 \\ \alpha & \zeta_7^3 - \zeta_7^4 & \zeta_7^5 - \zeta_7^2 \\ \zeta_7^3 - \zeta_7^4 & \zeta_7^5 - \zeta_7^2 & \alpha \end{pmatrix}$$

with  $\alpha := -1 - \zeta_7^5 - \zeta_7^4 - \zeta_7^3 - \zeta_7^2 - 2\zeta_7$

## Observations:

- $\mathbb{Q}[\zeta_7][x]^G$  is generated by:  $x_1x_2^3 + x_2x_3^3 + x_3x_1^3, 270x_3^2x_1^2x_2^2 - 54x_3^5x_1 - 54x_2^5x_3 - 54x_1^5x_2, f_3, f_4$  where  $f_3, f_4 \in \mathbb{Q}[x] \rightsquigarrow$   
Generating set of rational invariants!
- $\mathrm{Gal}(\mathbb{Q}[\zeta_7]/\mathbb{Q})$  acts on  $\Delta(G)$

Use this observation to give a precise formulation of this phenomena

- Let  $G$  be a finite group and  $\chi \in \text{Irr}(G)$  faithful

- Let  $G$  be a finite group and  $\chi \in \text{Irr}(G)$  faithful
- $K/\mathbb{Q}$  be GALOIS with  $\Gamma := \text{Gal}(K/\mathbb{Q})$

- Let  $G$  be a finite group and  $\chi \in \text{Irr}(G)$  faithful
- $K/\mathbb{Q}$  be GALOIS with  $\Gamma := \text{Gal}(K/\mathbb{Q})$
- $\Delta : G \rightarrow \text{GL}_n(K)$  a representation affording  $\chi$

- Let  $G$  be a finite group and  $\chi \in \text{Irr}(G)$  faithful
- $K/\mathbb{Q}$  be GALOIS with  $\Gamma := \text{Gal}(K/\mathbb{Q})$
- $\Delta : G \rightarrow \text{GL}_n(K)$  a representation affording  $\chi$

## Definition of a $K/\mathbb{Q}$ -form:

Define  $\Delta$  to be a  $K/\mathbb{Q}$ -form if there exists  $U \leq \text{Aut}(G)$  and an isomorphism  $\bar{\cdot} : U \rightarrow \Gamma$  such that

$$\Delta(u(g)) = \bar{u}(\Delta(g)) \text{ for every } g \in G$$

- Let  $G$  be a finite group and  $\chi \in \text{Irr}(G)$  faithful
- $K/\mathbb{Q}$  be GALOIS with  $\Gamma := \text{Gal}(K/\mathbb{Q})$
- $\Delta : G \rightarrow \text{GL}_n(K)$  a representation affording  $\chi$

## Definition of a $K/\mathbb{Q}$ -form:

Define  $\Delta$  to be a  $K/\mathbb{Q}$ -form if there exists  $U \leq \text{Aut}(G)$  and an isomorphism  $\bar{\cdot} : U \rightarrow \Gamma$  such that

$$\Delta(u(g)) = \bar{u}(\Delta(g)) \text{ for every } g \in G$$

- Trivial Example: A representation  $\Delta : G \rightarrow \text{GL}_n(\mathbb{Q})$  is a  $\mathbb{Q}/\mathbb{Q}$ -form with  $U = \langle 1 \rangle$ .

## Comparison of $K/\mathbb{Q}$ -forms:

Let  $\Delta$  and  $\Theta$  be  $K/\mathbb{Q}$ -forms then:  $\Delta \sim \Theta$  if and only if there exists  $Y \in \mathrm{GL}_n(\mathbb{Q})$  such that  $Y\Delta(g)Y^{-1} = \Theta(g)$  for all  $g \in G$

## Comparison of $K/\mathbb{Q}$ -forms:

Let  $\Delta$  and  $\Theta$  be  $K/\mathbb{Q}$ -forms then:  $\Delta \sim \Theta$  if and only if there exists  $Y \in \mathrm{GL}_n(\mathbb{Q})$  such that  $Y\Delta(g)Y^{-1} = \Theta(g)$  for all  $g \in G$

Connection with invariant theory:

## Comparison of $K/\mathbb{Q}$ -forms:

Let  $\Delta$  and  $\Theta$  be  $K/\mathbb{Q}$ -forms then:  $\Delta \sim \Theta$  if and only if there exists  $Y \in \mathrm{GL}_n(\mathbb{Q})$  such that  $Y\Delta(g)Y^{-1} = \Theta(g)$  for all  $g \in G$

Connection with invariant theory:

## Theorem:

$\Delta : G \rightarrow \mathrm{GL}_n(K)$  is a  $K/\mathbb{Q}$ -form if and only if  $K[x]^G$  is generated by polynomials with rational coefficients.

## Main question:

Given  $\chi \in \text{Irr}(G)$  and  $U \leq \text{Aut}(G)$  is there a GALOIS extension  $K$  of  $\mathbb{Q}$  such that there exists a  $K/\mathbb{Q}$ -form  $\Delta$  with this  $U$ ?

## Main question:

Given  $\chi \in \text{Irr}(G)$  and  $U \leq \text{Aut}(G)$  is there a GALOIS extension  $K$  of  $\mathbb{Q}$  such that there exists a  $K/\mathbb{Q}$ -form  $\Delta$  with this  $U$ ?

The following conditions are obviously necessary:

## Main question:

Given  $\chi \in \text{Irr}(G)$  and  $U \leq \text{Aut}(G)$  is there a GALOIS extension  $K$  of  $\mathbb{Q}$  such that there exists a  $K/\mathbb{Q}$ -form  $\Delta$  with this  $U$ ?

The following conditions are obviously necessary:

- There exists a GALOIS extension  $K/\mathbb{Q}$  with  $U$  as GALOIS group and it is possible to afford  $\chi$  as a  $K$ -representation.

## Main question:

Given  $\chi \in \text{Irr}(G)$  and  $U \leq \text{Aut}(G)$  is there a GALOIS extension  $K$  of  $\mathbb{Q}$  such that there exists a  $K/\mathbb{Q}$ -form  $\Delta$  with this  $U$ ?

The following conditions are obviously necessary:

- There exists a GALOIS extension  $K/\mathbb{Q}$  with  $U$  as GALOIS group and it is possible to afford  $\chi$  as a  $K$ -representation.
- $U$  acts transitively on  $\{\chi^\sigma \mid \sigma \in \text{Gal}(\mathbb{Q}(\chi)/\mathbb{Q})\}$  and  $\bar{u} \circ \chi = \chi \circ u$  for all  $u \in U$ .

# Existence a first approach

Assume we are given

- $\Delta : G \rightarrow \mathrm{GL}_n(K)$  a representation affording  $\chi$

# Existence a first approach

Assume we are given

- $\Delta : G \rightarrow \mathrm{GL}_n(K)$  a representation affording  $\chi$
- The necessary conditions on  $U$  and  $K$  of the previous slide are fulfilled. Especially  $\bar{u} \circ \chi = \chi \circ u$ .

Then it holds that:

# Existence a first approach

Assume we are given

- $\Delta : G \rightarrow \mathrm{GL}_n(K)$  a representation affording  $\chi$
- The necessary conditions on  $U$  and  $K$  of the previous slide are fulfilled. Especially  $\bar{u} \circ \chi = \chi \circ u$ .

Then it holds that:

- There exists  $X_u \in \mathrm{GL}_n(K)$  such that  $X_u \Delta(u(g)) X_u^{-1} = \bar{u}(\Delta(g))$

# Existence a first approach

Assume we are given

- $\Delta : G \rightarrow \mathrm{GL}_n(K)$  a representation affording  $\chi$
- The necessary conditions on  $U$  and  $K$  of the previous slide are fulfilled. Especially  $\bar{u} \circ \chi = \chi \circ u$ .

Then it holds that:

- There exists  $X_u \in \mathrm{GL}_n(K)$  such that  $X_u \Delta(u(g)) X_u^{-1} = \bar{u}(\Delta(g))$
- $\lambda_{s,t} = X_s^{-1} \bar{s}(X_t^{-1}) X_{st} \in Z^2(\Gamma, K^*)$  for  $s, t \in U$

# Existence a first approach

Assume we are given

- $\Delta : G \rightarrow \mathrm{GL}_n(K)$  a representation affording  $\chi$
- The necessary conditions on  $U$  and  $K$  of the previous slide are fulfilled. Especially  $\bar{u} \circ \chi = \chi \circ u$ .

Then it holds that:

- There exists  $X_u \in \mathrm{GL}_n(K)$  such that  $X_u \Delta(u(g)) X_u^{-1} = \bar{u}(\Delta(g))$
- $\lambda_{s,t} = X_s^{-1} \bar{s}(X_t^{-1}) X_{st} \in Z^2(\Gamma, K^*)$  for  $s, t \in U$

We get

Theorem:

There exists a  $K/\mathbb{Q}$ -form if and only if  $\lambda \sim 1 \in H^2(\Gamma, K^*)$ .

# Existence a first approach

## Problems:

- To calculate  $\lambda_{s,t}$  a representation  $\Delta : G \rightarrow \mathrm{GL}_n(K)$  is needed

# Existence a first approach

## Problems:

- To calculate  $\lambda_{s,t}$  a representation  $\Delta : G \rightarrow \mathrm{GL}_n(K)$  is needed
- Deciding the existence is almost the same as constructing the actual  $K/\mathbb{Q}$ -form

# Existence a first approach

## Problems:

- To calculate  $\lambda_{s,t}$  a representation  $\Delta : G \rightarrow \mathrm{GL}_n(K)$  is needed
- Deciding the existence is almost the same as constructing the actual  $K/\mathbb{Q}$ -form

## Goal:

# Existence a first approach

## Problems:

- To calculate  $\lambda_{s,t}$  a representation  $\Delta : G \rightarrow \mathrm{GL}_n(K)$  is needed
- Deciding the existence is almost the same as constructing the actual  $K/\mathbb{Q}$ -form

## Goal:

Decide existence of a  $K/\mathbb{Q}$ -form without constructing a representation  $\Delta : G \rightarrow \mathrm{GL}_n(K)$

# Existence a first approach

## Problems:

- To calculate  $\lambda_{s,t}$  a representation  $\Delta : G \rightarrow \mathrm{GL}_n(K)$  is needed
- Deciding the existence is almost the same as constructing the actual  $K/\mathbb{Q}$ -form

## Goal:

Decide existence of a  $K/\mathbb{Q}$ -form without constructing a representation  $\Delta : G \rightarrow \mathrm{GL}_n(K)$

Let  $e = \sum_{\sigma \in \mathrm{Gal}(\mathbb{Q}(\chi)/\mathbb{Q})} \sigma \circ e_\chi \in \mathbb{Q}G$  where  $e_\chi$  is the central primitive idempotent of  $\mathbb{C}[G]$  corresponding to  $\chi$ . Consider  $A := e\mathbb{Q}Ge$  as a simple  $\mathbb{Q}$ -algebra then:

# Existence a first approach

## Problems:

- To calculate  $\lambda_{s,t}$  a representation  $\Delta : G \rightarrow \mathrm{GL}_n(K)$  is needed
- Deciding the existence is almost the same as constructing the actual  $K/\mathbb{Q}$ -form

## Goal:

Decide existence of a  $K/\mathbb{Q}$ -form without constructing a representation  $\Delta : G \rightarrow \mathrm{GL}_n(K)$

Let  $e = \sum_{\sigma \in \mathrm{Gal}(\mathbb{Q}(\chi)/\mathbb{Q})} \sigma \circ e_\chi \in \mathbb{Q}G$  where  $e_\chi$  is the central primitive idempotent of  $\mathbb{C}[G]$  corresponding to  $\chi$ . Consider  $A := e\mathbb{Q}Ge$  as a simple  $\mathbb{Q}$ -algebra then:

- $U$  acts on  $\mathbb{Q}G$  as automorphisms by the  $\mathbb{Q}$ -linear extension of  ${}^u g := u(g)$

# Existence a first approach

## Problems:

- To calculate  $\lambda_{s,t}$  a representation  $\Delta : G \rightarrow \mathrm{GL}_n(K)$  is needed
- Deciding the existence is almost the same as constructing the actual  $K/\mathbb{Q}$ -form

## Goal:

Decide existence of a  $K/\mathbb{Q}$ -form without constructing a representation  $\Delta : G \rightarrow \mathrm{GL}_n(K)$

Let  $e = \sum_{\sigma \in \mathrm{Gal}(\mathbb{Q}(\chi)/\mathbb{Q})} \sigma \circ e_\chi \in \mathbb{Q}G$  where  $e_\chi$  is the central primitive idempotent of  $\mathbb{C}[G]$  corresponding to  $\chi$ . Consider  $A := e\mathbb{Q}Ge$  as a simple  $\mathbb{Q}$ -algebra then:

- $U$  acts on  $\mathbb{Q}G$  as automorphisms by the  $\mathbb{Q}$ -linear extension of  ${}^u g := u(g)$
- $U$  fixes  $e$  and so  $U$  acts on  $A$  as automorphisms

# Existence a first approach

## Problems:

- To calculate  $\lambda_{s,t}$  a representation  $\Delta : G \rightarrow \mathrm{GL}_n(K)$  is needed
- Deciding the existence is almost the same as constructing the actual  $K/\mathbb{Q}$ -form

## Goal:

Decide existence of a  $K/\mathbb{Q}$ -form without constructing a representation  $\Delta : G \rightarrow \mathrm{GL}_n(K)$

Let  $e = \sum_{\sigma \in \mathrm{Gal}(\mathbb{Q}(\chi)/\mathbb{Q})} \sigma \circ e_\chi \in \mathbb{Q}G$  where  $e_\chi$  is the central primitive idempotent of  $\mathbb{C}[G]$  corresponding to  $\chi$ . Consider  $A := e\mathbb{Q}Ge$  as a simple  $\mathbb{Q}$ -algebra then:

- $U$  acts on  $\mathbb{Q}G$  as automorphisms by the  $\mathbb{Q}$ -linear extension of  ${}^u g := u(g)$
- $U$  fixes  $e$  and so  $U$  acts on  $A$  as automorphisms
- For any representation  $\Delta : G \rightarrow \mathrm{GL}_n(K)$  affording  $\chi$  the action on  $A$  induces an action of  $U$  on  $\Delta(\mathbb{Q}G)$

So  $A$  is a simple  $\mathbb{Q}$ -algebra where  $U$  acts as automorphisms and  $A$  also is a central simple  $\mathbb{Q}(\chi)$ -algebra.

So  $A$  is a simple  $\mathbb{Q}$ -algebra where  $U$  acts as automorphisms and  $A$  also is a central simple  $\mathbb{Q}(\chi)$ -algebra.

Idea:

- Considering  $A$  as an element of the BRAUER group gives information about rationality questions of  $\chi$

So  $A$  is a simple  $\mathbb{Q}$ -algebra where  $U$  acts as automorphisms and  $A$  also is a central simple  $\mathbb{Q}(\chi)$ -algebra.

Idea:

- Considering  $A$  as an element of the BRAUER group gives information about rationality questions of  $\chi$
- Hope: Find a generalization of the BRAUER group which takes a group action into account and maybe answers the "rationality" question we are interested in.

The following definitions and theorems are due to TURULL

## Definition:

- A  **$G$ -algebra**  $A$  is a finite dimensional associative  $k$ -algebra where  $G$  acts as automorphisms

The following definitions and theorems are due to TURULL

## Definition:

- A  **$G$ -algebra**  $A$  is a finite dimensional associative  $k$ -algebra where  $G$  acts as automorphisms
- $A$  is a **simple  $G$ -algebra** if it has only trivial 2-sided  $G$ -invariant ideals

The following definitions and theorems are due to TURULL

## Definition:

- A  **$G$ -algebra**  $A$  is a finite dimensional associative  $k$ -algebra where  $G$  acts as automorphisms
- $A$  is a **simple  $G$ -algebra** if it has only trivial 2-sided  $G$ -invariant ideals
- $A$  is called **central** if  $\text{Fix}_{C(A)}(G) = k$

The following definitions and theorems are due to TURULL

## Definition:

- A  **$G$ -algebra**  $A$  is a finite dimensional associative  $k$ -algebra where  $G$  acts as automorphisms
- $A$  is a **simple  $G$ -algebra** if it has only trivial 2-sided  $G$ -invariant ideals
- $A$  is called **central** if  $\text{Fix}_{C(A)}(G) = k$
- Two  $G$ -algebras are called isomorphic if there exist a  $G$ -equivariant  $k$ -algebra isomorphism  $\varphi$

The following definitions and theorems are due to TURULL

## Definition:

- A  **$G$ -algebra**  $A$  is a finite dimensional associative  $k$ -algebra where  $G$  acts as automorphisms
- $A$  is a **simple  $G$ -algebra** if it has only trivial 2-sided  $G$ -invariant ideals
- $A$  is called **central** if  $\text{Fix}_{C(A)}(G) = k$
- Two  $G$ -algebras are called isomorphic if there exist a  $G$ -equivariant  $k$ -algebra isomorphism  $\varphi$
- A  $G$ -algebra  $A$  is called **trivial** if there exist a  $kG$ -module  $M$  and  $A \cong_G \text{End}_k(M)$  where the  $G$ -structure on  $\text{End}_k(M)$  is given by conjugation

The following definitions and theorems are due to TURULL

## Definition:

- A  **$G$ -algebra**  $A$  is a finite dimensional associative  $k$ -algebra where  $G$  acts as automorphisms
- $A$  is a **simple  $G$ -algebra** if it has only trivial 2-sided  $G$ -invariant ideals
- $A$  is called **central** if  $\text{Fix}_{C(A)}(G) = k$
- Two  $G$ -algebras are called isomorphic if there exist a  $G$ -equivariant  $k$ -algebra isomorphism  $\varphi$
- A  $G$ -algebra  $A$  is called **trivial** if there exist a  $kG$ -module  $M$  and  $A \cong_G \text{End}_k(M)$  where the  $G$ -structure on  $\text{End}_k(M)$  is given by conjugation
- We call two  $G$ -algebras  $A$  and  $B$  **equivalent** if there exists trivial  $G$ -algebras  $E_1$  and  $E_2$  such that:  $A \otimes_k E_1 \cong_G B \otimes_k E_2$ . We simply write  $A \sim_G B$

The following theorem defines our generalization of the BRAUER group

The following theorem defines our generalization of the BRAUER group

Theorem: [Turull 09]

Let  $F$  be a  $G$ -field with  $F^G = k$ . We define  $\text{BrCliff}(G, F)$  as set of all equivalence classes of central simple  $G$ -algebras such that the centres are  $G$ -isomorphic to  $F$ . Then  $\text{BrCliff}(G, F)$  is an abelian group with the following group structure:

$$\text{BrCliff}(G, F) \times \text{BrCliff}(G, F) \rightarrow \text{BrCliff}(G, F)$$

$$([A], [B]) \mapsto [A \otimes_F B]$$

where  $G$  acts diagonally on the tensor product

## Remark and notation:

- Forgetting about the  $G$ -action gives a homomorphism  
 $\text{BrCliff}(G, F) \rightarrow \text{Br}(F)^{\text{Gal}(F/k)}$

## Remark and notation:

- Forgetting about the  $G$ -action gives a homomorphism  $\text{BrCliff}(G, F) \rightarrow \text{Br}(F)^{\text{Gal}(F/k)}$
- Denote the kernel of this homomorphism by  $\text{FMBrCliff}(G, F)$

## Remark and notation:

- Forgetting about the  $G$ -action gives a homomorphism  $\text{BrCliff}(G, F) \rightarrow \text{Br}(F)^{\text{Gal}(F/k)}$
- Denote the kernel of this homomorphism by  $\text{FMBrCliff}(G, F)$
- For a central simple  $G$ -algebra  $A$  denote by  $[A]$  it's element in  $\text{BrCliff}(G, F)$

## Remark and notation:

- Forgetting about the  $G$ -action gives a homomorphism  $\text{BrCliff}(G, F) \rightarrow \text{Br}(F)^{\text{Gal}(F/k)}$
- Denote the kernel of this homomorphism by  $\text{FMBrCliff}(G, F)$
- For a central simple  $G$ -algebra  $A$  denote by  $[A]$  it's element in  $\text{BrCliff}(G, F)$

## Theorem: [Turull 09]

$$\text{FMBrCliff}(G, F) \cong H^2(G, F^*)$$

## Remark and notation:

- Forgetting about the  $G$ -action gives a homomorphism  $\text{BrCliff}(G, F) \rightarrow \text{Br}(F)^{\text{Gal}(F/k)}$
- Denote the kernel of this homomorphism by  $\text{FMBrCliff}(G, F)$
- For a central simple  $G$ -algebra  $A$  denote by  $[A]$  it's element in  $\text{BrCliff}(G, F)$

## Theorem: [Turull 09]

$$\text{FMBrCliff}(G, F) \cong H^2(G, F^*)$$

## Assumption: $m_\chi = 1$

## Remark and notation:

- Forgetting about the  $G$ -action gives a homomorphism  $\text{BrCliff}(G, F) \rightarrow \text{Br}(F)^{\text{Gal}(F/k)}$
- Denote the kernel of this homomorphism by  $\text{FMBrCliff}(G, F)$
- For a central simple  $G$ -algebra  $A$  denote by  $[A]$  its element in  $\text{BrCliff}(G, F)$

## Theorem: [Turull 09]

$$\text{FMBrCliff}(G, F) \cong H^2(G, F^*)$$

## Assumption: $m_\chi = 1$

Then  $[A] := [e\mathbb{Q}Ge] \in \text{FMBrCliff}(U, \mathbb{Q}(\chi))$  where the  $U$  action was defined earlier.

# Main Theorem

Using the last theorem we get:

# Main Theorem

Using the last theorem we get:

## Theorem:

Let  $G$  be a finite group,  $\chi \in \text{Irr}(G)$  faithful with SCHUR index one over  $\mathbb{Q}$ . Given  $U \leq \text{Aut}(G)$  and a GALOIS field  $K$  such that the previous conditions are fulfilled, then there exists a  $K/\mathbb{Q}$ -form if and only if  $[A] = [e\mathbb{Q}Ge] = [1] \in \text{BrCliff}(U, \mathbb{Q}(\chi))$ .

# Main Theorem

Using the last theorem we get:

## Theorem:

Let  $G$  be a finite group,  $\chi \in \text{Irr}(G)$  faithful with SCHUR index one over  $\mathbb{Q}$ . Given  $U \leq \text{Aut}(G)$  and a GALOIS field  $K$  such that the previous conditions are fulfilled, then there exists a  $K/\mathbb{Q}$ -form if and only if  $[A] = [e\mathbb{Q}Ge] = [1] \in \text{BrCliff}(U, \mathbb{Q}(\chi))$ .

From this we directly get the following corollary.

# Main Theorem

Using the last theorem we get:

## Theorem:

Let  $G$  be a finite group,  $\chi \in \text{Irr}(G)$  faithful with SCHUR index one over  $\mathbb{Q}$ . Given  $U \leq \text{Aut}(G)$  and a GALOIS field  $K$  such that the previous conditions are fulfilled, then there exists a  $K/\mathbb{Q}$ -form if and only if  $[A] = [e\mathbb{Q}Ge] = [1] \in \text{BrCliff}(U, \mathbb{Q}(\chi))$ .

From this we directly get the following corollary.

## Corollary:

With the assumptions of the last theorem: There exists a  $K/\mathbb{Q}$ -form if and only if the irreducible  $\mathbb{Q}G$  module  $M$  corresponding to the character  $\sum_{\sigma \in \text{Gal}(\mathbb{Q}(\chi)/\mathbb{Q})} \sigma \circ \chi$  extends to an irreducible  $\mathbb{Q}G \rtimes U$ -module.

# Conclusions

Upshot:

## Upshot:

- If  $m_\chi = 1$  then deciding existence of a  $K/\mathbb{Q}$ -form can in principle be done without constructing a representation

## Upshot:

- If  $m_\chi = 1$  then deciding existence of a  $K/\mathbb{Q}$ -form can in principle be done without constructing a representation
- Formulation could be used to prove existence for the groups  $\mathrm{PSL}(2, q)$

# Conclusions

## Upshot:

- If  $m_\chi = 1$  then deciding existence of a  $K/\mathbb{Q}$ -form can in principle be done without constructing a representation
- Formulation could be used to prove existence for the groups  $\mathrm{PSL}(2, q)$

## Open Problems

# Conclusions

## Upshot:

- If  $m_\chi = 1$  then deciding existence of a  $K/\mathbb{Q}$ -form can in principle be done without constructing a representation
- Formulation could be used to prove existence for the groups  $\mathrm{PSL}(2, q)$

## Open Problems

- Link the existence problem in general to properties of the  $U$ -algebra  $e\mathbb{Q}Ge \rightsquigarrow$  in principle avoiding calculating a representation

## Upshot:

- If  $m_\chi = 1$  then deciding existence of a  $K/\mathbb{Q}$ -form can in principle be done without constructing a representation
- Formulation could be used to prove existence for the groups  $\mathrm{PSL}(2, q)$

## Open Problems

- Link the existence problem in general to properties of the  $U$ -algebra  $e\mathbb{Q}Ge \rightsquigarrow$  in principle avoiding calculating a representation
- Describe  $\mathrm{BrCliff}(G, K)$  in terms of GALOIS cohomology

## Upshot:

- If  $m_\chi = 1$  then deciding existence of a  $K/\mathbb{Q}$ -form can in principle be done without constructing a representation
- Formulation could be used to prove existence for the groups  $\mathrm{PSL}(2, q)$

## Open Problems

- Link the existence problem in general to properties of the  $U$ -algebra  $e\mathbb{Q}Ge \rightsquigarrow$  in principle avoiding calculating a representation
- Describe  $\mathrm{BrCliff}(G, K)$  in terms of GALOIS cohomology
- Find and implement algorithms in MAGMA or GAP