

# ON THE LIE ALGEBRA OF SKEW-SYMMETRIC ELEMENTS OF AN ENVELOPING ALGEBRA

Salvatore Siciliano

Dipartimento di Matematica “E. De Giorgi”  
Università del Salento  
salvatore.siciliano@unisalento.it

Aachen, March 22-26, 2010

Let  $A$  be an algebra with involution  $*$ . Denote by

$$A^+ := \{x \in A \mid x^* = x\}$$

the set of *symmetric elements* of  $A$  under  $*$ , and by

$$A^- := \{x \in A \mid x^* = -x\}$$

the set of *skew-symmetric elements* of  $A$  under  $*$ .

Theorem (Amitsur, 1968)

*Let  $A$  be an algebra with involution. If  $A^+$  or  $A^-$  satisfies a polynomial identity, then so does  $A$ .*

Let  $A$  be an algebra with involution  $*$ . Denote by

$$A^+ := \{x \in A \mid x^* = x\}$$

the set of *symmetric elements* of  $A$  under  $*$ , and by

$$A^- := \{x \in A \mid x^* = -x\}$$

the set of *skew-symmetric elements* of  $A$  under  $*$ .

Theorem (Amitsur, 1968)

*Let  $A$  be an algebra with involution. If  $A^+$  or  $A^-$  satisfies a polynomial identity, then so does  $A$ .*

Let  $A$  be an algebra with involution  $*$ . Denote by

$$A^+ := \{x \in A \mid x^* = x\}$$

the set of *symmetric elements* of  $A$  under  $*$ , and by

$$A^- := \{x \in A \mid x^* = -x\}$$

the set of *skew-symmetric elements* of  $A$  under  $*$ .

Theorem (Amitsur, 1968)

*Let  $A$  be an algebra with involution. If  $A^+$  or  $A^-$  satisfies a polynomial identity, then so does  $A$ .*

Let  $A$  be an algebra with involution  $*$ . Denote by

$$A^+ := \{x \in A \mid x^* = x\}$$

the set of *symmetric elements* of  $A$  under  $*$ , and by

$$A^- := \{x \in A \mid x^* = -x\}$$

the set of *skew-symmetric elements* of  $A$  under  $*$ .

Theorem (Amitsur, 1968)

*Let  $A$  be an algebra with involution. If  $A^+$  or  $A^-$  satisfies a polynomial identity, then so does  $A$ .*

Let  $A$  be an algebra with involution  $*$ . Denote by

$$A^+ := \{x \in A \mid x^* = x\}$$

the set of *symmetric elements* of  $A$  under  $*$ , and by

$$A^- := \{x \in A \mid x^* = -x\}$$

the set of *skew-symmetric elements* of  $A$  under  $*$ .

Theorem (Amitsur, 1968)

*Let  $A$  be an algebra with involution. If  $A^+$  or  $A^-$  satisfies a polynomial identity, then so does  $A$ .*

Let  $A$  be an algebra with involution  $*$ . Denote by

$$A^+ := \{x \in A \mid x^* = x\}$$

the set of *symmetric elements* of  $A$  under  $*$ , and by

$$A^- := \{x \in A \mid x^* = -x\}$$

the set of *skew-symmetric elements* of  $A$  under  $*$ .

### Theorem (Amitsur, 1968)

*Let  $A$  be an algebra with involution. If  $A^+$  or  $A^-$  satisfies a polynomial identity, then so does  $A$ .*

Let  $A$  be an algebra with involution  $*$ . Denote by

$$A^+ := \{x \in A \mid x^* = x\}$$

the set of *symmetric elements* of  $A$  under  $*$ , and by

$$A^- := \{x \in A \mid x^* = -x\}$$

the set of *skew-symmetric elements* of  $A$  under  $*$ .

### Theorem (Amitsur, 1968)

*Let  $A$  be an algebra with involution. If  $A^+$  or  $A^-$  satisfies a polynomial identity, then so does  $A$ .*

Let  $A$  be an algebra with involution.

Note that  $A^-$  has a structure of a Lie algebra induced by the Lie product of  $A$  (defined by  $[x, y] := xy - yx$  for every  $x, y \in A$ ).

Let  $F$  be a field and  $G$  a group.

We consider the group algebra  $\mathbb{F}G$  endowed with the *canonical involution* (given by linear extension of the inversion map  $g \mapsto g^{-1}$  of  $G$ ).

Let  $A$  be an algebra with involution.

Note that  $A^-$  has a structure of a Lie algebra induced by the Lie product of  $A$  (defined by  $[x, y] := xy - yx$  for every  $x, y \in A$ ).

Let  $F$  be a field and  $G$  a group.

We consider the group algebra  $\mathbb{F}G$  endowed with the *canonical involution* (given by linear extension of the inversion map  $g \mapsto g^{-1}$  of  $G$ ).

Let  $A$  be an algebra with involution.

Note that  $A^-$  has a structure of a Lie algebra induced by the Lie product of  $A$  (defined by  $[x, y] := xy - yx$  for every  $x, y \in A$ ).

Let  $F$  be a field and  $G$  a group.

We consider the group algebra  $\mathbb{F}G$  endowed with the *canonical involution* (given by linear extension of the inversion map  $g \mapsto g^{-1}$  of  $G$ ).

Let  $A$  be an algebra with involution.

Note that  $A^-$  has a structure of a Lie algebra induced by the Lie product of  $A$  (defined by  $[x, y] := xy - yx$  for every  $x, y \in A$ ).

Let  $F$  be a field and  $G$  a group.

We consider the group algebra  $\mathbb{F}G$  endowed with the *canonical involution* (given by linear extension of the inversion map  $g \mapsto g^{-1}$  of  $G$ ).

Let  $p$  be a prime. We say that group  $G$  is  $p$ -abelian if  $G'$  is a finite  $p$ -group. By a 0-abelian group we mean simply an abelian group.

Theorem (Giambruno-Sehgal, 1993, 2006)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and let  $G$  be a group. Then  $\mathbb{F}G^-$  is nilpotent if and only if one the following conditions holds:

- 1)  $G$  has a nilpotent  $p$ -abelian normal subgroup  $H$  with  $(G \setminus H)^2 = \{1\}$ ;
- 2)  $G$  has an elementary abelian 2-subgroup of index 2;
- 3) the set  $P$  of  $p$ -elements of  $G$  forms a finite normal subgroup of  $G$  and  $G/P$  is an elementary abelian 2-group.

Let  $p$  be a prime. We say that group  $G$  is  $p$ -abelian if  $G'$  is a finite  $p$ -group. By a 0-abelian group we mean simply an abelian group.

Theorem (Giambruno-Sehgal, 1993, 2006)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and let  $G$  be a group. Then  $\mathbb{F}G^-$  is nilpotent if and only if one the following conditions holds:

- 1)  $G$  has a nilpotent  $p$ -abelian normal subgroup  $H$  with  $(G \setminus H)^2 = \{1\}$ ;
- 2)  $G$  has an elementary abelian 2-subgroup of index 2;
- 3) the set  $P$  of  $p$ -elements of  $G$  forms a finite normal subgroup of  $G$  and  $G/P$  is an elementary abelian 2-group.

Let  $p$  be a prime. We say that group  $G$  is  $p$ -abelian if  $G'$  is a finite  $p$ -group. By a 0-abelian group we mean simply an abelian group.

Theorem (Giambruno-Sehgal, 1993, 2006)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and let  $G$  be a group. Then  $\mathbb{F}G^-$  is nilpotent if and only if one the following conditions holds:

- 1)  $G$  has a nilpotent  $p$ -abelian normal subgroup  $H$  with  $(G \setminus H)^2 = \{1\}$ ;
- 2)  $G$  has an elementary abelian 2-subgroup of index 2;
- 3) the set  $P$  of  $p$ -elements of  $G$  forms a finite normal subgroup of  $G$  and  $G/P$  is an elementary abelian 2-group.

Let  $p$  be a prime. We say that group  $G$  is  $p$ -abelian if  $G'$  is a finite  $p$ -group. By a 0-abelian group we mean simply an abelian group.

### Theorem (Giambruno-Sehgal, 1993, 2006)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and let  $G$  be a group. Then  $\mathbb{F}G^-$  is nilpotent if and only if one the following conditions holds:

- 1)  $G$  has a nilpotent  $p$ -abelian normal subgroup  $H$  with  $(G \setminus H)^2 = \{1\}$ ;
- 2)  $G$  has an elementary abelian 2-subgroup of index 2;
- 3) the set  $P$  of  $p$ -elements of  $G$  forms a finite normal subgroup of  $G$  and  $G/P$  is an elementary abelian 2-group.

Let  $p$  be a prime. We say that group  $G$  is  $p$ -abelian if  $G'$  is a finite  $p$ -group. By a 0-abelian group we mean simply an abelian group.

### Theorem (Giambruno-Sehgal, 1993, 2006)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and let  $G$  be a group. Then  $\mathbb{F}G^-$  is nilpotent if and only if one the following conditions holds:

- 1)  $G$  has a nilpotent  $p$ -abelian normal subgroup  $H$  with  $(G \setminus H)^2 = \{1\}$ ;
- 2)  $G$  has an elementary abelian 2-subgroup of index 2;
- 3) the set  $P$  of  $p$ -elements of  $G$  forms a finite normal subgroup of  $G$  and  $G/P$  is an elementary abelian 2-group.

Let  $p$  be a prime. We say that group  $G$  is  $p$ -abelian if  $G'$  is a finite  $p$ -group. By a 0-abelian group we mean simply an abelian group.

### Theorem (Giambruno-Sehgal, 1993, 2006)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and let  $G$  be a group. Then  $\mathbb{F}G^-$  is nilpotent if and only if one the following conditions holds:

- 1)  $G$  has a nilpotent  $p$ -abelian normal subgroup  $H$  with  $(G \setminus H)^2 = \{1\}$ ;
- 2)  $G$  has an elementary abelian 2-subgroup of index 2;
- 3) the set  $P$  of  $p$ -elements of  $G$  forms a finite normal subgroup of  $G$  and  $G/P$  is an elementary abelian 2-group.

Let  $p$  be a prime. We say that group  $G$  is  $p$ -abelian if  $G'$  is a finite  $p$ -group. By a 0-abelian group we mean simply an abelian group.

### Theorem (Giambruno-Sehgal, 1993, 2006)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and let  $G$  be a group. Then  $\mathbb{F}G^-$  is nilpotent if and only if one the following conditions holds:

- 1)  $G$  has a nilpotent  $p$ -abelian normal subgroup  $H$  with  $(G \setminus H)^2 = \{1\}$ ;
- 2)  $G$  has an elementary abelian 2-subgroup of index 2;
- 3) the set  $P$  of  $p$ -elements of  $G$  forms a finite normal subgroup of  $G$  and  $G/P$  is an elementary abelian 2-group.

Let  $p$  be a prime. We say that group  $G$  is  $p$ -abelian if  $G'$  is a finite  $p$ -group. By a 0-abelian group we mean simply an abelian group.

### Theorem (Giambruno-Sehgal, 1993, 2006)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and let  $G$  be a group. Then  $\mathbb{F}G^-$  is nilpotent if and only if one the following conditions holds:

- 1)  $G$  has a nilpotent  $p$ -abelian normal subgroup  $H$  with  $(G \setminus H)^2 = \{1\}$ ;
- 2)  $G$  has an elementary abelian 2-subgroup of index 2;
- 3) the set  $P$  of  $p$ -elements of  $G$  forms a finite normal subgroup of  $G$  and  $G/P$  is an elementary abelian 2-group.

## Theorem (Lee, 2000)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and let  $G$  be a group with no 2-elements. Then the following conditions are equivalent:

- 1)  $\mathbb{F}G^-$  is  $n$ -Engel for some  $n$ ;
- 2)  $\mathbb{F}G$  is Lie  $m$ -Engel for some  $m$ ;
- 3) either
  - (i)  $p = 0$  and  $G$  is abelian or
  - (ii)  $p > 2$ ,  $G$  is nilpotent and there exists a normal subgroup  $A$  of  $G$  such that  $G/A$  and  $A'$  are both finite  $p$ -groups.

## Theorem (Lee, 2000)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and let  $G$  be a group with no 2-elements. Then the following conditions are equivalent:

- 1)  $\mathbb{F}G^-$  is  $n$ -Engel for some  $n$ ;
- 2)  $\mathbb{F}G$  is Lie  $m$ -Engel for some  $m$ ;
- 3) either
  - (i)  $p = 0$  and  $G$  is abelian or
  - (ii)  $p > 2$ ,  $G$  is nilpotent and there exists a normal subgroup  $A$  of  $G$  such that  $G/A$  and  $A'$  are both finite  $p$ -groups.

## Theorem (Lee, 2000)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and let  $G$  be a group with no 2-elements. Then the following conditions are equivalent:

- 1)  $\mathbb{F}G^-$  is  $n$ -Engel for some  $n$ ;
- 2)  $\mathbb{F}G$  is Lie  $m$ -Engel for some  $m$ ;
- 3) either
  - (i)  $p = 0$  and  $G$  is abelian or
  - (ii)  $p > 2$ ,  $G$  is nilpotent and there exists a normal subgroup  $A$  of  $G$  such that  $G/A$  and  $A'$  are both finite  $p$ -groups.

### Theorem (Lee, 2000)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and let  $G$  be a group with no 2-elements. Then the following conditions are equivalent:

- 1)  $\mathbb{F}G^-$  is  $n$ -Engel for some  $n$ ;
- 2)  $\mathbb{F}G$  is Lie  $m$ -Engel for some  $m$ ;
- 3) either
  - (i)  $p = 0$  and  $G$  is abelian or
  - (ii)  $p > 2$ ,  $G$  is nilpotent and there exists a normal subgroup  $A$  of  $G$  such that  $G/A$  and  $A'$  are both finite  $p$ -groups.

### Theorem (Lee, 2000)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and let  $G$  be a group with no 2-elements. Then the following conditions are equivalent:

- 1)  $\mathbb{F}G^-$  is  $n$ -Engel for some  $n$ ;
- 2)  $\mathbb{F}G$  is Lie  $m$ -Engel for some  $m$ ;
- 3) either
  - (i)  $p = 0$  and  $G$  is abelian or
  - (ii)  $p > 2$ ,  $G$  is nilpotent and there exists a normal subgroup  $A$  of  $G$  such that  $G/A$  and  $A'$  are both finite  $p$ -groups.

## Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}G$  be the group algebra of a group  $G$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$ . If the set  $P$  of the  $p$ -elements of  $G$  is finite, then  $FG^-$  is solvable if and only if  $P$  is a normal subgroup of  $G$  and one of the following conditions occurs:

- 1)  $G/P$  is abelian;
- 2)  $G/P = A \rtimes \langle x \rangle$ , where  $A$  is abelian,  $o(x) = 2$ , and  $x$  acts dihedrally upon  $A$ ;
- 3)  $G/P$  contains an elementary abelian 2-subgroup of index 2.

## Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and  $G$  a group. If the set  $P$  of  $p$ -elements of  $G$  contains an infinite subgroup of bounded exponent, and  $G$  contains no nontrivial elements of order dividing  $p^2 - 1$ , then the following conditions are equivalent:

- 1)  $FG^-$  is solvable;
- 2)  $FG$  is Lie solvable;
- 3)  $G'$  is a finite  $p$ -group.

### Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}G$  be the group algebra of a group  $G$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$ . If the set  $P$  of the  $p$ -elements of  $G$  is finite, then  $FG^-$  is solvable if and only if  $P$  is a normal subgroup of  $G$  and one of the following conditions occurs:

- 1)  $G/P$  is abelian;
- 2)  $G/P = A \rtimes \langle x \rangle$ , where  $A$  is abelian,  $o(x) = 2$ , and  $x$  acts dihedrally upon  $A$ ;
- 3)  $G/P$  contains an elementary abelian 2-subgroup of index 2.

### Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and  $G$  a group. If the set  $P$  of  $p$ -elements of  $G$  contains an infinite subgroup of bounded exponent, and  $G$  contains no nontrivial elements of order dividing  $p^2 - 1$ , then the following conditions are equivalent:

- 1)  $FG^-$  is solvable;
- 2)  $FG$  is Lie solvable;
- 3)  $G'$  is a finite  $p$ -group.

## Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}G$  be the group algebra of a group  $G$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$ . If the set  $P$  of the  $p$ -elements of  $G$  is finite, then  $FG^-$  is solvable if and only if  $P$  is a normal subgroup of  $G$  and one of the following conditions occurs:

- 1)  $G/P$  is abelian;
- 2)  $G/P = A \rtimes \langle x \rangle$ , where  $A$  is abelian,  $o(x) = 2$ , and  $x$  acts dihedrally upon  $A$ ;
- 3)  $G/P$  contains an elementary abelian 2-subgroup of index 2.

## Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and  $G$  a group. If the set  $P$  of  $p$ -elements of  $G$  contains an infinite subgroup of bounded exponent, and  $G$  contains no nontrivial elements of order dividing  $p^2 - 1$ , then the following conditions are equivalent:

- 1)  $FG^-$  is solvable;
- 2)  $FG$  is Lie solvable;
- 3)  $G'$  is a finite  $p$ -group.

### Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}G$  be the group algebra of a group  $G$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$ . If the set  $P$  of the  $p$ -elements of  $G$  is finite, then  $FG^-$  is solvable if and only if  $P$  is a normal subgroup of  $G$  and one of the following conditions occurs:

- 1)  $G/P$  is abelian;
- 2)  $G/P = A \rtimes \langle x \rangle$ , where  $A$  is abelian,  $o(x) = 2$ , and  $x$  acts dihedrally upon  $A$ ;
- 3)  $G/P$  contains an elementary abelian 2-subgroup of index 2.

### Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and  $G$  a group. If the set  $P$  of  $p$ -elements of  $G$  contains an infinite subgroup of bounded exponent, and  $G$  contains no nontrivial elements of order dividing  $p^2 - 1$ , then the following conditions are equivalent:

- 1)  $FG^-$  is solvable;
- 2)  $FG$  is Lie solvable;
- 3)  $G'$  is a finite  $p$ -group.

### Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}G$  be the group algebra of a group  $G$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$ . If the set  $P$  of the  $p$ -elements of  $G$  is finite, then  $FG^-$  is solvable if and only if  $P$  is a normal subgroup of  $G$  and one of the following conditions occurs:

- 1)  $G/P$  is abelian;
- 2)  $G/P = A \rtimes \langle x \rangle$ , where  $A$  is abelian,  $o(x) = 2$ , and  $x$  acts dihedrally upon  $A$ ;
- 3)  $G/P$  contains an elementary abelian 2-subgroup of index 2.

### Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and  $G$  a group. If the set  $P$  of  $p$ -elements of  $G$  contains an infinite subgroup of bounded exponent, and  $G$  contains no nontrivial elements of order dividing  $p^2 - 1$ , then the following conditions are equivalent:

- 1)  $FG^-$  is solvable;
- 2)  $FG$  is Lie solvable;
- 3)  $G'$  is a finite  $p$ -group.

### Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}G$  be the group algebra of a group  $G$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$ . If the set  $P$  of the  $p$ -elements of  $G$  is finite, then  $FG^-$  is solvable if and only if  $P$  is a normal subgroup of  $G$  and one of the following conditions occurs:

- 1)  $G/P$  is abelian;
- 2)  $G/P = A \rtimes \langle x \rangle$ , where  $A$  is abelian,  $o(x) = 2$ , and  $x$  acts dihedrally upon  $A$ ;
- 3)  $G/P$  contains an elementary abelian 2-subgroup of index 2.

### Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and  $G$  a group. If the set  $P$  of  $p$ -elements of  $G$  contains an infinite subgroup of bounded exponent, and  $G$  contains no nontrivial elements of order dividing  $p^2 - 1$ , then the following conditions are equivalent:

- 1)  $FG^-$  is solvable;
- 2)  $FG$  is Lie solvable;
- 3)  $G'$  is a finite  $p$ -group.

### Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}G$  be the group algebra of a group  $G$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$ . If the set  $P$  of the  $p$ -elements of  $G$  is finite, then  $FG^-$  is solvable if and only if  $P$  is a normal subgroup of  $G$  and one of the following conditions occurs:

- 1)  $G/P$  is abelian;
- 2)  $G/P = A \rtimes \langle x \rangle$ , where  $A$  is abelian,  $o(x) = 2$ , and  $x$  acts dihedrally upon  $A$ ;
- 3)  $G/P$  contains an elementary abelian 2-subgroup of index 2.

### Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and  $G$  a group. If the set  $P$  of  $p$ -elements of  $G$  contains an infinite subgroup of bounded exponent, and  $G$  contains no nontrivial elements of order dividing  $p^2 - 1$ , then the following conditions are equivalent:

- 1)  $FG^-$  is solvable;
- 2)  $FG$  is Lie solvable;
- 3)  $G'$  is a finite  $p$ -group.

### Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}G$  be the group algebra of a group  $G$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$ . If the set  $P$  of the  $p$ -elements of  $G$  is finite, then  $FG^-$  is solvable if and only if  $P$  is a normal subgroup of  $G$  and one of the following conditions occurs:

- 1)  $G/P$  is abelian;
- 2)  $G/P = A \rtimes \langle x \rangle$ , where  $A$  is abelian,  $o(x) = 2$ , and  $x$  acts dihedrally upon  $A$ ;
- 3)  $G/P$  contains an elementary abelian 2-subgroup of index 2.

### Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and  $G$  a group. If the set  $P$  of  $p$ -elements of  $G$  contains an infinite subgroup of bounded exponent, and  $G$  contains no nontrivial elements of order dividing  $p^2 - 1$ , then the following conditions are equivalent:

- 1)  $FG^-$  is solvable;
- 2)  $FG$  is Lie solvable;
- 3)  $G'$  is a finite  $p$ -group.

### Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}G$  be the group algebra of a group  $G$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$ . If the set  $P$  of the  $p$ -elements of  $G$  is finite, then  $FG^-$  is solvable if and only if  $P$  is a normal subgroup of  $G$  and one of the following conditions occurs:

- 1)  $G/P$  is abelian;
- 2)  $G/P = A \rtimes \langle x \rangle$ , where  $A$  is abelian,  $o(x) = 2$ , and  $x$  acts dihedrally upon  $A$ ;
- 3)  $G/P$  contains an elementary abelian 2-subgroup of index 2.

### Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and  $G$  a group. If the set  $P$  of  $p$ -elements of  $G$  contains an infinite subgroup of bounded exponent, and  $G$  contains no nontrivial elements of order dividing  $p^2 - 1$ , then the following conditions are equivalent:

- 1)  $FG^-$  is solvable;
- 2)  $FG$  is Lie solvable;
- 3)  $G'$  is a finite  $p$ -group.

### Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}G$  be the group algebra of a group  $G$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$ . If the set  $P$  of the  $p$ -elements of  $G$  is finite, then  $FG^-$  is solvable if and only if  $P$  is a normal subgroup of  $G$  and one of the following conditions occurs:

- 1)  $G/P$  is abelian;
- 2)  $G/P = A \rtimes \langle x \rangle$ , where  $A$  is abelian,  $o(x) = 2$ , and  $x$  acts dihedrally upon  $A$ ;
- 3)  $G/P$  contains an elementary abelian 2-subgroup of index 2.

### Theorem (Lee-Sehgal-Spinelli, 2009)

Let  $\mathbb{F}$  be a field of characteristic  $p \neq 2$ , and  $G$  a group. If the set  $P$  of  $p$ -elements of  $G$  contains an infinite subgroup of bounded exponent, and  $G$  contains no nontrivial elements of order dividing  $p^2 - 1$ , then the following conditions are equivalent:

- 1)  $FG^-$  is solvable;
- 2)  $FG$  is Lie solvable;
- 3)  $G'$  is a finite  $p$ -group.

For a restricted Lie algebra  $L$  over a field  $\mathbb{F}$  of characteristic  $p > 0$ , we denote by  $u(L)$  the restricted enveloping algebra of  $L$ .

### Theorem (Riley-Shalev, 1993)

*Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 0$ .*

- 1)  $u(L)$  is Lie nilpotent if and only if  $L$  is nilpotent and  $L'$  is finite-dimensional and  $p$ -nilpotent;*
- 2)  $u(L)$  is Lie  $n$ -Engel for some  $n$  if and only if  $L$  is nilpotent,  $L'$  is  $p$ -nilpotent, and  $L$  contains a restricted ideal  $I$  such that  $L/I$  and  $I'$  are finite-dimensional;*
- 3) If  $p > 2$ , then  $u(L)$  is Lie solvable if and only if  $L'$  is finite-dimensional and  $p$ -nilpotent.*

For a restricted Lie algebra  $L$  over a field  $\mathbb{F}$  of characteristic  $p > 0$ , we denote by  $u(L)$  the restricted enveloping algebra of  $L$ .

### Theorem (Riley-Shalev, 1993)

*Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 0$ .*

- 1)  $u(L)$  is Lie nilpotent if and only if  $L$  is nilpotent and  $L'$  is finite-dimensional and  $p$ -nilpotent;*
- 2)  $u(L)$  is Lie  $n$ -Engel for some  $n$  if and only if  $L$  is nilpotent,  $L'$  is  $p$ -nilpotent, and  $L$  contains a restricted ideal  $I$  such that  $L/I$  and  $I'$  are finite-dimensional;*
- 3) If  $p > 2$ , then  $u(L)$  is Lie solvable if and only if  $L'$  is finite-dimensional and  $p$ -nilpotent.*

For a restricted Lie algebra  $L$  over a field  $\mathbb{F}$  of characteristic  $p > 0$ , we denote by  $u(L)$  the restricted enveloping algebra of  $L$ .

### Theorem (Riley-Shalev, 1993)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 0$ .

- 1)  $u(L)$  is Lie nilpotent if and only if  $L$  is nilpotent and  $L'$  is finite-dimensional and  $p$ -nilpotent;
- 2)  $u(L)$  is Lie  $n$ -Engel for some  $n$  if and only if  $L$  is nilpotent,  $L'$  is  $p$ -nilpotent, and  $L$  contains a restricted ideal  $I$  such that  $L/I$  and  $I'$  are finite-dimensional;
- 3) If  $p > 2$ , then  $u(L)$  is Lie solvable if and only if  $L'$  is finite-dimensional and  $p$ -nilpotent.

For a restricted Lie algebra  $L$  over a field  $\mathbb{F}$  of characteristic  $p > 0$ , we denote by  $u(L)$  the restricted enveloping algebra of  $L$ .

### Theorem (Riley-Shalev, 1993)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 0$ .

- 1)  $u(L)$  is Lie nilpotent if and only if  $L$  is nilpotent and  $L'$  is finite-dimensional and  $p$ -nilpotent;
- 2)  $u(L)$  is Lie  $n$ -Engel for some  $n$  if and only if  $L$  is nilpotent,  $L'$  is  $p$ -nilpotent, and  $L$  contains a restricted ideal  $I$  such that  $L/I$  and  $I'$  are finite-dimensional;
- 3) If  $p > 2$ , then  $u(L)$  is Lie solvable if and only if  $L'$  is finite-dimensional and  $p$ -nilpotent.

For a restricted Lie algebra  $L$  over a field  $\mathbb{F}$  of characteristic  $p > 0$ , we denote by  $u(L)$  the restricted enveloping algebra of  $L$ .

### Theorem (Riley-Shalev, 1993)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 0$ .

- 1)  $u(L)$  is Lie nilpotent if and only if  $L$  is nilpotent and  $L'$  is finite-dimensional and  $p$ -nilpotent;
- 2)  $u(L)$  is Lie  $n$ -Engel for some  $n$  if and only if  $L$  is nilpotent,  $L'$  is  $p$ -nilpotent, and  $L$  contains a restricted ideal  $I$  such that  $L/I$  and  $I'$  are finite-dimensional;
- 3) If  $p > 2$ , then  $u(L)$  is Lie solvable if and only if  $L'$  is finite-dimensional and  $p$ -nilpotent.

For a restricted Lie algebra  $L$  over a field  $\mathbb{F}$  of characteristic  $p > 0$ , we denote by  $u(L)$  the restricted enveloping algebra of  $L$ .

### Theorem (Riley-Shalev, 1993)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 0$ .

- 1)  $u(L)$  is Lie nilpotent if and only if  $L$  is nilpotent and  $L'$  is finite-dimensional and  $p$ -nilpotent;
- 2)  $u(L)$  is Lie  $n$ -Engel for some  $n$  if and only if  $L$  is nilpotent,  $L'$  is  $p$ -nilpotent, and  $L$  contains a restricted ideal  $I$  such that  $L/I$  and  $I'$  are finite-dimensional;
- 3) If  $p > 2$ , then  $u(L)$  is Lie solvable if and only if  $L'$  is finite-dimensional and  $p$ -nilpotent.

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 0$ . We consider  $u(L)$  endowed with the *principal involution*, that is, the unique  $\mathbb{F}$ -antiautomorphism  $\top$  of  $u(L)$  such that  $x^\top = -x$  for every  $x$  in  $L$ .

If  $\mathcal{L}$  is an arbitrary Lie algebra, the *principal involution* of the (ordinary) universal enveloping algebra  $U(\mathcal{L})$  of  $\mathcal{L}$  is defined in a similar way.

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 0$ . We consider  $u(L)$  endowed with the *principal involution*, that is, the unique  $\mathbb{F}$ -antiautomorphism  $\top$  of  $u(L)$  such that  $x^\top = -x$  for every  $x$  in  $L$ .

If  $\mathcal{L}$  is an arbitrary Lie algebra, the *principal involution* of the (ordinary) universal enveloping algebra  $U(\mathcal{L})$  of  $\mathcal{L}$  is defined in a similar way.

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 0$ . We consider  $u(L)$  endowed with the *principal involution*, that is, the unique  $\mathbb{F}$ -antiautomorphism  $\top$  of  $u(L)$  such that  $x^\top = -x$  for every  $x$  in  $L$ .

If  $\mathcal{L}$  is an arbitrary Lie algebra, the *principal involution* of the (ordinary) universal enveloping algebra  $U(\mathcal{L})$  of  $\mathcal{L}$  is defined in a similar way.

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 0$ . We consider  $u(L)$  endowed with the *principal involution*, that is, the unique  $\mathbb{F}$ -antiautomorphism  $\top$  of  $u(L)$  such that  $x^\top = -x$  for every  $x$  in  $L$ .

If  $\mathcal{L}$  is an arbitrary Lie algebra, the *principal involution* of the (ordinary) universal enveloping algebra  $U(\mathcal{L})$  of  $\mathcal{L}$  is defined in a similar way.

## Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is solvable;
- 2)  $u(L)$  is Lie solvable;
- 3)  $L'$  is finite-dimensional and  $p$ -nilpotent.

## Example

Let  $L$  be the restricted Lie algebra over a field  $\mathbb{F}$  of characteristic 2 with a basis  $\{x_1, x_2, y, z\}$  such that  $[x_1, x_2] = z$ ,  $[x_1, y] = x_1$ ,  $[x_2, y] = x_2$ ,  $z \in Z(L)$ ,  $x_1^{[2]} = x_2^{[2]} = 0$ ,  $y^{[2]} = y$  and  $z^{[p]} = z$ . Then  $u(L)^-$  is solvable, but  $u(L)$  is not Lie solvable.

## Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is solvable;
- 2)  $u(L)$  is Lie solvable;
- 3)  $L'$  is finite-dimensional and  $p$ -nilpotent.

## Example

Let  $L$  be the restricted Lie algebra over a field  $\mathbb{F}$  of characteristic 2 with a basis  $\{x_1, x_2, y, z\}$  such that  $[x_1, x_2] = z$ ,  $[x_1, y] = x_1$ ,  $[x_2, y] = x_2$ ,  $z \in Z(L)$ ,  $x_1^{[2]} = x_2^{[2]} = 0$ ,  $y^{[2]} = y$  and  $z^{[p]} = z$ . Then  $u(L)^-$  is solvable, but  $u(L)$  is not Lie solvable.

## Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is solvable;
- 2)  $u(L)$  is Lie solvable;
- 3)  $L'$  is finite-dimensional and  $p$ -nilpotent.

## Example

Let  $L$  be the restricted Lie algebra over a field  $\mathbb{F}$  of characteristic 2 with a basis  $\{x_1, x_2, y, z\}$  such that  $[x_1, x_2] = z$ ,  $[x_1, y] = x_1$ ,  $[x_2, y] = x_2$ ,  $z \in Z(L)$ ,  $x_1^{[2]} = x_2^{[2]} = 0$ ,  $y^{[2]} = y$  and  $z^{[p]} = z$ . Then  $u(L)^-$  is solvable, but  $u(L)$  is not Lie solvable.

## Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is solvable;
- 2)  $u(L)$  is Lie solvable;
- 3)  $L'$  is finite-dimensional and  $p$ -nilpotent.

## Example

Let  $L$  be the restricted Lie algebra over a field  $\mathbb{F}$  of characteristic 2 with a basis  $\{x_1, x_2, y, z\}$  such that  $[x_1, x_2] = z$ ,  $[x_1, y] = x_1$ ,  $[x_2, y] = x_2$ ,  $z \in Z(L)$ ,  $x_1^{[2]} = x_2^{[2]} = 0$ ,  $y^{[2]} = y$  and  $z^{[p]} = z$ . Then  $u(L)^-$  is solvable, but  $u(L)$  is not Lie solvable.

## Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is solvable;
- 2)  $u(L)$  is Lie solvable;
- 3)  $L'$  is finite-dimensional and  $p$ -nilpotent.

## Example

Let  $L$  be the restricted Lie algebra over a field  $\mathbb{F}$  of characteristic 2 with a basis  $\{x_1, x_2, y, z\}$  such that  $[x_1, x_2] = z$ ,  $[x_1, y] = x_1$ ,  $[x_2, y] = x_2$ ,  $z \in Z(L)$ ,  $x_1^{[2]} = x_2^{[2]} = 0$ ,  $y^{[2]} = y$  and  $z^{[p]} = z$ . Then  $u(L)^-$  is solvable, but  $u(L)$  is not Lie solvable.

## Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is solvable;
- 2)  $u(L)$  is Lie solvable;
- 3)  $L'$  is finite-dimensional and  $p$ -nilpotent.

## Example

Let  $L$  be the restricted Lie algebra over a field  $\mathbb{F}$  of characteristic 2 with a basis  $\{x_1, x_2, y, z\}$  such that  $[x_1, x_2] = z$ ,  $[x_1, y] = x_1$ ,  $[x_2, y] = x_2$ ,  $z \in Z(L)$ ,  $x_1^{[2]} = x_2^{[2]} = 0$ ,  $y^{[2]} = y$  and  $z^{[p]} = z$ . Then  $u(L)^-$  is solvable, but  $u(L)$  is not Lie solvable.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is solvable;
- 2)  $u(L)$  is Lie solvable;
- 3)  $L'$  is finite-dimensional and  $p$ -nilpotent.

### Example

Let  $L$  be the restricted Lie algebra over a field  $\mathbb{F}$  of characteristic 2 with a basis  $\{x_1, x_2, y, z\}$  such that  $[x_1, x_2] = z$ ,  $[x_1, y] = x_1$ ,  $[x_2, y] = x_2$ ,  $z \in Z(L)$ ,  $x_1^{[2]} = x_2^{[2]} = 0$ ,  $y^{[2]} = y$  and  $z^{[p]} = z$ . Then  $u(L)^-$  is solvable, but  $u(L)$  is not Lie solvable.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is solvable;
- 2)  $u(L)$  is Lie solvable;
- 3)  $L'$  is finite-dimensional and  $p$ -nilpotent.

### Example

Let  $L$  be the restricted Lie algebra over a field  $\mathbb{F}$  of characteristic 2 with a basis  $\{x_1, x_2, y, z\}$  such that  $[x_1, x_2] = z$ ,  $[x_1, y] = x_1$ ,  $[x_2, y] = x_2$ ,  $z \in Z(L)$ ,  $x_1^{[2]} = x_2^{[2]} = 0$ ,  $y^{[2]} = y$  and  $z^{[p]} = z$ . Then  $u(L)^-$  is solvable, but  $u(L)$  is not Lie solvable.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is solvable;
- 2)  $u(L)$  is Lie solvable;
- 3)  $L'$  is finite-dimensional and  $p$ -nilpotent.

### Example

Let  $L$  be the restricted Lie algebra over a field  $\mathbb{F}$  of characteristic 2 with a basis  $\{x_1, x_2, y, z\}$  such that  $[x_1, x_2] = z$ ,  $[x_1, y] = x_1$ ,  $[x_2, y] = x_2$ ,  $z \in Z(L)$ ,  $x_1^{[2]} = x_2^{[2]} = 0$ ,  $y^{[2]} = y$  and  $z^{[p]} = z$ . Then  $u(L)^-$  is solvable, but  $u(L)$  is not Lie solvable.

## Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is  $n$ -Engel for some  $n$ ;
- 2)  $u(L)$  is Lie  $m$ -Engel for some  $m$ ;
- 3)  $L$  is nilpotent,  $L'$  is  $p$ -nilpotent, and  $L$  contains a restricted ideal  $I$  such that  $L/I$  and  $I'$  are finite-dimensional.

## Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is nilpotent;
- 2)  $u(L)$  is Lie nilpotent;
- 3)  $L$  is nilpotent and  $L'$  is finite-dimensional and  $p$ -nilpotent.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is  $n$ -Engel for some  $n$ ;
- 2)  $u(L)$  is Lie  $m$ -Engel for some  $m$ ;
- 3)  $L$  is nilpotent,  $L'$  is  $p$ -nilpotent, and  $L$  contains a restricted ideal  $I$  such that  $L/I$  and  $I'$  are finite-dimensional.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is nilpotent;
- 2)  $u(L)$  is Lie nilpotent;
- 3)  $L$  is nilpotent and  $L'$  is finite-dimensional and  $p$ -nilpotent.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is  $n$ -Engel for some  $n$ ;
- 2)  $u(L)$  is Lie  $m$ -Engel for some  $m$ ;
- 3)  $L$  is nilpotent,  $L'$  is  $p$ -nilpotent, and  $L$  contains a restricted ideal  $I$  such that  $L/I$  and  $I'$  are finite-dimensional.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is nilpotent;
- 2)  $u(L)$  is Lie nilpotent;
- 3)  $L$  is nilpotent and  $L'$  is finite-dimensional and  $p$ -nilpotent.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is  $n$ -Engel for some  $n$ ;
- 2)  $u(L)$  is Lie  $m$ -Engel for some  $m$ ;
- 3)  $L$  is nilpotent,  $L'$  is  $p$ -nilpotent, and  $L$  contains a restricted ideal  $I$  such that  $L/I$  and  $I'$  are finite-dimensional.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is nilpotent;
- 2)  $u(L)$  is Lie nilpotent;
- 3)  $L$  is nilpotent and  $L'$  is finite-dimensional and  $p$ -nilpotent.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is  $n$ -Engel for some  $n$ ;
- 2)  $u(L)$  is Lie  $m$ -Engel for some  $m$ ;
- 3)  $L$  is nilpotent,  $L'$  is  $p$ -nilpotent, and  $L$  contains a restricted ideal  $I$  such that  $L/I$  and  $I'$  are finite-dimensional.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is nilpotent;
- 2)  $u(L)$  is Lie nilpotent;
- 3)  $L$  is nilpotent and  $L'$  is finite-dimensional and  $p$ -nilpotent.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is  $n$ -Engel for some  $n$ ;
- 2)  $u(L)$  is Lie  $m$ -Engel for some  $m$ ;
- 3)  $L$  is nilpotent,  $L'$  is  $p$ -nilpotent, and  $L$  contains a restricted ideal  $I$  such that  $L/I$  and  $I'$  are finite-dimensional.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is nilpotent;
- 2)  $u(L)$  is Lie nilpotent;
- 3)  $L$  is nilpotent and  $L'$  is finite-dimensional and  $p$ -nilpotent.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is  $n$ -Engel for some  $n$ ;
- 2)  $u(L)$  is Lie  $m$ -Engel for some  $m$ ;
- 3)  $L$  is nilpotent,  $L'$  is  $p$ -nilpotent, and  $L$  contains a restricted ideal  $I$  such that  $L/I$  and  $I'$  are finite-dimensional.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is nilpotent;
- 2)  $u(L)$  is Lie nilpotent;
- 3)  $L$  is nilpotent and  $L'$  is finite-dimensional and  $p$ -nilpotent.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is  $n$ -Engel for some  $n$ ;
- 2)  $u(L)$  is Lie  $m$ -Engel for some  $m$ ;
- 3)  $L$  is nilpotent,  $L'$  is  $p$ -nilpotent, and  $L$  contains a restricted ideal  $I$  such that  $L/I$  and  $I'$  are finite-dimensional.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is nilpotent;
- 2)  $u(L)$  is Lie nilpotent;
- 3)  $L$  is nilpotent and  $L'$  is finite-dimensional and  $p$ -nilpotent.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is  $n$ -Engel for some  $n$ ;
- 2)  $u(L)$  is Lie  $m$ -Engel for some  $m$ ;
- 3)  $L$  is nilpotent,  $L'$  is  $p$ -nilpotent, and  $L$  contains a restricted ideal  $I$  such that  $L/I$  and  $I'$  are finite-dimensional.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is nilpotent;
- 2)  $u(L)$  is Lie nilpotent;
- 3)  $L$  is nilpotent and  $L'$  is finite-dimensional and  $p$ -nilpotent.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is  $n$ -Engel for some  $n$ ;
- 2)  $u(L)$  is Lie  $m$ -Engel for some  $m$ ;
- 3)  $L$  is nilpotent,  $L'$  is  $p$ -nilpotent, and  $L$  contains a restricted ideal  $I$  such that  $L/I$  and  $I'$  are finite-dimensional.

### Theorem (S., 2010)

Let  $L$  be a restricted Lie algebra over a field  $\mathbb{F}$  of characteristic  $p > 2$ . Then the following conditions are equivalent:

- 1)  $u(L)^-$  is nilpotent;
- 2)  $u(L)$  is Lie nilpotent;
- 3)  $L$  is nilpotent and  $L'$  is finite-dimensional and  $p$ -nilpotent.

## Example

Let  $L$  be the restricted Lie algebra over a field  $\mathbb{F}$  of characteristic 2 with a basis  $\{x, y, z\}$  such that  $[x, y] = z$ ,  $z \in Z(L)$ ,  $x^{[2]} = y^{[2]} = 0$  and  $z^{[2]} = z$ . Then  $u(L)^-$  is nilpotent, but  $u(L)$  cannot satisfy any Engel condition.

In odd characteristic,  $u(L)^-$  is solvable, nilpotent or  $n$ -Engel for some  $n$  if and only if so is  $u(L)$  as a Lie algebra.

## Example

Let  $L$  be the restricted Lie algebra over a field  $\mathbb{F}$  of characteristic 2 with a basis  $\{x, y, z\}$  such that  $[x, y] = z$ ,  $z \in Z(L)$ ,  $x^{[2]} = y^{[2]} = 0$  and  $z^{[2]} = z$ . Then  $u(L)^-$  is nilpotent, but  $u(L)$  cannot satisfy any Engel condition.

In odd characteristic,  $u(L)^-$  is solvable, nilpotent or  $n$ -Engel for some  $n$  if and only if so is  $u(L)$  as a Lie algebra.

## Example

Let  $L$  be the restricted Lie algebra over a field  $\mathbb{F}$  of characteristic 2 with a basis  $\{x, y, z\}$  such that  $[x, y] = z$ ,  $z \in Z(L)$ ,  $x^{[2]} = y^{[2]} = 0$  and  $z^{[2]} = z$ . Then  $u(L)^-$  is nilpotent, but  $u(L)$  cannot satisfy any Engel condition.

In odd characteristic,  $u(L)^-$  is solvable, nilpotent or  $n$ -Engel for some  $n$  if and only if so is  $u(L)$  as a Lie algebra.

## Example

Let  $L$  be the restricted Lie algebra over a field  $\mathbb{F}$  of characteristic 2 with a basis  $\{x, y, z\}$  such that  $[x, y] = z$ ,  $z \in Z(L)$ ,  $x^{[2]} = y^{[2]} = 0$  and  $z^{[2]} = z$ . Then  $u(L)^-$  is nilpotent, but  $u(L)$  cannot satisfy any Engel condition.

In odd characteristic,  $u(L)^-$  is solvable, nilpotent or  $n$ -Engel for some  $n$  if and only if so is  $u(L)$  as a Lie algebra.

## Example

Let  $L$  be the restricted Lie algebra over a field  $\mathbb{F}$  of characteristic 2 with a basis  $\{x, y, z\}$  such that  $[x, y] = z$ ,  $z \in Z(L)$ ,  $x^{[2]} = y^{[2]} = 0$  and  $z^{[2]} = z$ . Then  $u(L)^-$  is nilpotent, but  $u(L)$  cannot satisfy any Engel condition.

In odd characteristic,  $u(L)^-$  is solvable, nilpotent or  $n$ -Engel for some  $n$  if and only if so is  $u(L)$  as a Lie algebra.

## Example

Let  $L$  be the restricted Lie algebra over a field  $\mathbb{F}$  of characteristic 2 with a basis  $\{x, y, z\}$  such that  $[x, y] = z$ ,  $z \in Z(L)$ ,  $x^{[2]} = y^{[2]} = 0$  and  $z^{[2]} = z$ . Then  $u(L)^-$  is nilpotent, but  $u(L)$  cannot satisfy any Engel condition.

In odd characteristic,  $u(L)^-$  is solvable, nilpotent or  $n$ -Engel for some  $n$  if and only if so is  $u(L)$  as a Lie algebra.

## Example

Let  $L$  be the restricted Lie algebra over a field  $\mathbb{F}$  of characteristic 2 with a basis  $\{x, y, z\}$  such that  $[x, y] = z$ ,  $z \in Z(L)$ ,  $x^{[2]} = y^{[2]} = 0$  and  $z^{[2]} = z$ . Then  $u(L)^-$  is nilpotent, but  $u(L)$  cannot satisfy any Engel condition.

In odd characteristic,  $u(L)^-$  is solvable, nilpotent or  $n$ -Engel for some  $n$  if and only if so is  $u(L)$  as a Lie algebra.

## Example

Let  $L$  be the restricted Lie algebra over a field  $\mathbb{F}$  of characteristic 2 with a basis  $\{x, y, z\}$  such that  $[x, y] = z$ ,  $z \in Z(L)$ ,  $x^{[2]} = y^{[2]} = 0$  and  $z^{[2]} = z$ . Then  $u(L)^-$  is nilpotent, but  $u(L)$  cannot satisfy any Engel condition.

In odd characteristic,  $u(L)^-$  is solvable, nilpotent or  $n$ -Engel for some  $n$  if and only if so is  $u(L)$  as a Lie algebra.

## Example

Let  $L$  be the restricted Lie algebra over a field  $\mathbb{F}$  of characteristic 2 with a basis  $\{x, y, z\}$  such that  $[x, y] = z$ ,  $z \in Z(L)$ ,  $x^{[2]} = y^{[2]} = 0$  and  $z^{[2]} = z$ . Then  $u(L)^-$  is nilpotent, but  $u(L)$  cannot satisfy any Engel condition.

In odd characteristic,  $u(L)^-$  is solvable, nilpotent or  $n$ -Engel for some  $n$  if and only if so is  $u(L)$  as a Lie algebra.

Now, consider any Lie algebra  $L$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$  and assume that  $U(L)^-$  is solvable or  $n$ -Engel for some  $n$ .

If  $p = 0$ , as  $U(L)$  satisfies a polynomial identity (by Amitsur's Theorem), then  $L$  is necessarily abelian (by Latyšev's Theorem).

Suppose then  $p > 2$ . Put

$$\hat{L} := \sum_{k \geq 0} L^{p^k} \subseteq U(L)$$

where  $L^{p^k} = \langle l^{p^k} \mid l \in L \rangle_{\mathbb{F}}$ . Then  $\hat{L}$  is a restricted Lie algebra with  $h^{[p]} = h^p$  for all  $h \in \hat{L}$ , and  $U(L) = u(\hat{L})$ . Therefore  $\hat{L}'$  is  $p$ -nilpotent and then, since  $U(L)$  is a domain, we conclude that  $L' = 0$ .

Now, consider any Lie algebra  $L$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$  and assume that  $U(L)^-$  is solvable or  $n$ -Engel for some  $n$ .

If  $p = 0$ , as  $U(L)$  satisfies a polynomial identity (by Amitsur's Theorem), then  $L$  is necessarily abelian (by Latyšev's Theorem).

Suppose then  $p > 2$ . Put

$$\hat{L} := \sum_{k \geq 0} L^{p^k} \subseteq U(L)$$

where  $L^{p^k} = \langle l^{p^k} \mid l \in L \rangle_{\mathbb{F}}$ . Then  $\hat{L}$  is a restricted Lie algebra with  $h^{[p]} = h^p$  for all  $h \in \hat{L}$ , and  $U(L) = u(\hat{L})$ . Therefore  $\hat{L}'$  is  $p$ -nilpotent and then, since  $U(L)$  is a domain, we conclude that  $L' = 0$ .

Now, consider any Lie algebra  $L$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$  and assume that  $U(L)^-$  is solvable or  $n$ -Engel for some  $n$ .

If  $p = 0$ , as  $U(L)$  satisfies a polynomial identity (by Amitsur's Theorem), then  $L$  is necessarily abelian (by Latyšev's Theorem).

Suppose then  $p > 2$ . Put

$$\hat{L} := \sum_{k \geq 0} L^{p^k} \subseteq U(L)$$

where  $L^{p^k} = \langle l^{p^k} \mid l \in L \rangle_{\mathbb{F}}$ . Then  $\hat{L}$  is a restricted Lie algebra with  $h^{[p]} = h^p$  for all  $h \in \hat{L}$ , and  $U(L) = u(\hat{L})$ . Therefore  $\hat{L}'$  is  $p$ -nilpotent and then, since  $U(L)$  is a domain, we conclude that  $L' = 0$ .

Now, consider any Lie algebra  $L$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$  and assume that  $U(L)^-$  is solvable or  $n$ -Engel for some  $n$ .

If  $p = 0$ , as  $U(L)$  satisfies a polynomial identity (by Amitsur's Theorem), then  $L$  is necessarily abelian (by Latyšev's Theorem).

Suppose then  $p > 2$ . Put

$$\hat{L} := \sum_{k \geq 0} L^{p^k} \subseteq U(L)$$

where  $L^{p^k} = \langle l^{p^k} \mid l \in L \rangle_{\mathbb{F}}$ . Then  $\hat{L}$  is a restricted Lie algebra with  $h^{[p]} = h^p$  for all  $h \in \hat{L}$ , and  $U(L) = u(\hat{L})$ . Therefore  $\hat{L}'$  is  $p$ -nilpotent and then, since  $U(L)$  is a domain, we conclude that  $L' = 0$ .

Now, consider any Lie algebra  $L$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$  and assume that  $U(L)^-$  is solvable or  $n$ -Engel for some  $n$ .

If  $p = 0$ , as  $U(L)$  satisfies a polynomial identity (by Amitsur's Theorem), then  $L$  is necessarily abelian (by Latyšev's Theorem).

Suppose then  $p > 2$ . Put

$$\hat{L} := \sum_{k \geq 0} L^{p^k} \subseteq U(L)$$

where  $L^{p^k} = \langle I^{p^k} \mid I \in L \rangle_{\mathbb{F}}$ . Then  $\hat{L}$  is a restricted Lie algebra with  $h^{[p]} = h^p$  for all  $h \in \hat{L}$ , and  $U(L) = u(\hat{L})$ . Therefore  $\hat{L}'$  is  $p$ -nilpotent and then, since  $U(L)$  is a domain, we conclude that  $L' = 0$ .

Now, consider any Lie algebra  $L$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$  and assume that  $U(L)^-$  is solvable or  $n$ -Engel for some  $n$ .

If  $p = 0$ , as  $U(L)$  satisfies a polynomial identity (by Amitsur's Theorem), then  $L$  is necessarily abelian (by Latyšev's Theorem).

Suppose then  $p > 2$ . Put

$$\hat{L} := \sum_{k \geq 0} L^{p^k} \subseteq U(L)$$

where  $L^{p^k} = \langle l^{p^k} \mid l \in L \rangle_{\mathbb{F}}$ . Then  $\hat{L}$  is a restricted Lie algebra with  $h^{[p]} = h^p$  for all  $h \in \hat{L}$ , and  $U(L) = u(\hat{L})$ . Therefore  $\hat{L}'$  is  $p$ -nilpotent and then, since  $U(L)$  is a domain, we conclude that  $L' = 0$ .

Now, consider any Lie algebra  $L$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$  and assume that  $U(L)^-$  is solvable or  $n$ -Engel for some  $n$ .

If  $p = 0$ , as  $U(L)$  satisfies a polynomial identity (by Amitsur's Theorem), then  $L$  is necessarily abelian (by Latyšev's Theorem).

Suppose then  $p > 2$ . Put

$$\hat{L} := \sum_{k \geq 0} L^{p^k} \subseteq U(L)$$

where  $L^{p^k} = \langle l^{p^k} \mid l \in L \rangle_{\mathbb{F}}$ . Then  $\hat{L}$  is a restricted Lie algebra with  $h^{[p]} = h^p$  for all  $h \in \hat{L}$ , and  $U(L) = u(\hat{L})$ . Therefore  $\hat{L}'$  is  $p$ -nilpotent and then, since  $U(L)$  is a domain, we conclude that  $L' = 0$ .

Now, consider any Lie algebra  $L$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$  and assume that  $U(L)^-$  is solvable or  $n$ -Engel for some  $n$ .

If  $p = 0$ , as  $U(L)$  satisfies a polynomial identity (by Amitsur's Theorem), then  $L$  is necessarily abelian (by Latyšev's Theorem).

Suppose then  $p > 2$ . Put

$$\hat{L} := \sum_{k \geq 0} L^{p^k} \subseteq U(L)$$

where  $L^{p^k} = \langle l^{p^k} \mid l \in L \rangle_{\mathbb{F}}$ . Then  $\hat{L}$  is a restricted Lie algebra with  $h^{[p]} = h^p$  for all  $h \in \hat{L}$ , and  $U(L) = u(\hat{L})$ . Therefore  $\hat{L}'$  is  $p$ -nilpotent and then, since  $U(L)$  is a domain, we conclude that  $L' = 0$ .

Now, consider any Lie algebra  $L$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$  and assume that  $U(L)^-$  is solvable or  $n$ -Engel for some  $n$ .

If  $p = 0$ , as  $U(L)$  satisfies a polynomial identity (by Amitsur's Theorem), then  $L$  is necessarily abelian (by Latyšev's Theorem).

Suppose then  $p > 2$ . Put

$$\hat{L} := \sum_{k \geq 0} L^{p^k} \subseteq U(L)$$

where  $L^{p^k} = \langle l^{p^k} \mid l \in L \rangle_{\mathbb{F}}$ . Then  $\hat{L}$  is a restricted Lie algebra with  $h^{[p]} = h^p$  for all  $h \in \hat{L}$ , and  $U(L) = u(\hat{L})$ . Therefore  $\hat{L}'$  is  $p$ -nilpotent and then, since  $U(L)$  is a domain, we conclude that  $L' = 0$ .

Now, consider any Lie algebra  $L$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$  and assume that  $U(L)^-$  is solvable or  $n$ -Engel for some  $n$ .

If  $p = 0$ , as  $U(L)$  satisfies a polynomial identity (by Amitsur's Theorem), then  $L$  is necessarily abelian (by Latyšev's Theorem).

Suppose then  $p > 2$ . Put

$$\hat{L} := \sum_{k \geq 0} L^{p^k} \subseteq U(L)$$

where  $L^{p^k} = \langle l^{p^k} \mid l \in L \rangle_{\mathbb{F}}$ . Then  $\hat{L}$  is a restricted Lie algebra with  $h^{[p]} = h^p$  for all  $h \in \hat{L}$ , and  $U(L) = u(\hat{L})$ . Therefore  $\hat{L}'$  is  $p$ -nilpotent and then, since  $U(L)$  is a domain, we conclude that  $L' = 0$ .

Now, consider any Lie algebra  $L$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$  and assume that  $U(L)^-$  is solvable or  $n$ -Engel for some  $n$ .

If  $p = 0$ , as  $U(L)$  satisfies a polynomial identity (by Amitsur's Theorem), then  $L$  is necessarily abelian (by Latyšev's Theorem).

Suppose then  $p > 2$ . Put

$$\hat{L} := \sum_{k \geq 0} L^{p^k} \subseteq U(L)$$

where  $L^{p^k} = \langle l^{p^k} \mid l \in L \rangle_{\mathbb{F}}$ . Then  $\hat{L}$  is a restricted Lie algebra with  $h^{[p]} = h^p$  for all  $h \in \hat{L}$ , and  $U(L) = u(\hat{L})$ . Therefore  $\hat{L}'$  is  $p$ -nilpotent and then, since  $U(L)$  is a domain, we conclude that  $L' = 0$ .

Now, consider any Lie algebra  $L$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$  and assume that  $U(L)^-$  is solvable or  $n$ -Engel for some  $n$ .

If  $p = 0$ , as  $U(L)$  satisfies a polynomial identity (by Amitsur's Theorem), then  $L$  is necessarily abelian (by Latyšev's Theorem).

Suppose then  $p > 2$ . Put

$$\hat{L} := \sum_{k \geq 0} L^{p^k} \subseteq U(L)$$

where  $L^{p^k} = \langle l^{p^k} \mid l \in L \rangle_{\mathbb{F}}$ . Then  $\hat{L}$  is a restricted Lie algebra with  $h^{[p]} = h^p$  for all  $h \in \hat{L}$ , and  $U(L) = u(\hat{L})$ . Therefore  $\hat{L}'$  is  $p$ -nilpotent and then, since  $U(L)$  is a domain, we conclude that  $L' = 0$ .

Now, consider any Lie algebra  $L$  over a field  $\mathbb{F}$  of characteristic  $p \neq 2$  and assume that  $U(L)^-$  is solvable or  $n$ -Engel for some  $n$ .

If  $p = 0$ , as  $U(L)$  satisfies a polynomial identity (by Amitsur's Theorem), then  $L$  is necessarily abelian (by Latyšev's Theorem).

Suppose then  $p > 2$ . Put

$$\hat{L} := \sum_{k \geq 0} L^{p^k} \subseteq U(L)$$

where  $L^{p^k} = \langle l^{p^k} \mid l \in L \rangle_{\mathbb{F}}$ . Then  $\hat{L}$  is a restricted Lie algebra with  $h^{[p]} = h^p$  for all  $h \in \hat{L}$ , and  $U(L) = u(\hat{L})$ . Therefore  $\hat{L}'$  is  $p$ -nilpotent and then, since  $U(L)$  is a domain, we conclude that  $L' = 0$ .

## Corollary (S., 2010)

*Let  $L$  be a Lie algebra over a field  $\mathbb{F}$  of characteristic  $p \neq 2$ . Then  $U(L)^-$  is solvable or  $n$ -Engel for some  $n$  if and only if  $L$  is abelian.*

### Corollary (S., 2010)

*Let  $L$  be a Lie algebra over a field  $\mathbb{F}$  of characteristic  $p \neq 2$ . Then  $U(L)^-$  is solvable or  $n$ -Engel for some  $n$  if and only if  $L$  is abelian.*

### Corollary (S., 2010)

*Let  $L$  be a Lie algebra over a field  $\mathbb{F}$  of characteristic  $p \neq 2$ . Then  $U(L)^-$  is solvable or  $n$ -Engel for some  $n$  if and only if  $L$  is abelian.*