

Nilpotency of group ring units symmetric with respect to an involution

Ernesto Spinelli

Università del Salento
Dipartimento di Matematica "E. De Giorgi"
Joint work with Gregory T. Lee and Sudarshan K. Sehgal

Aachen, March 22-26, 2010
Arithmetic of group rings and related structures

Outline

- 1 Involutions in group rings
 - Classical and K -linear involutions
 - Symmetric and skew-symmetric elements
 - Symmetric and unitary units
- 2 Main questions
 - Main question for symmetric and skew-symmetric elements
 - Main question for symmetric units
- 3 Group identities for symmetric units
 - The classical involution: an overview
 - K -linear involutions: the main results

Canonical and K -linear involutions

Let G be a group endowed with an involution \star .

Let us consider the K -linear extension of \star to KG by setting

$$\left(\sum_{g \in G} a_g g \right)^\star := \sum_{g \in G} a_g g^\star.$$

This extension, which we denote again by \star , is an involution of KG which fixes the ground field K elementwise.

As well-known, any group G has a natural involution which is given by the map $\ast : g \mapsto g^{-1}$.

Definition

Let KG be the group algebra of a group G over a field K . If G is endowed with an involution \ast , its linear extension to the group algebra KG is called a *K -linear involution* of KG . In particular, if $\star = \ast$ the induced involution is called *the classical involution*.

Canonical and K -linear involutions

Let G be a group endowed with an involution \star .

Let us consider the K -linear extension of \star to KG by setting

$$\left(\sum_{g \in G} a_g g \right)^\star := \sum_{g \in G} a_g g^\star.$$

This extension, which we denote again by \star , is an involution of KG which fixes the ground field K elementwise.

As well-known, any group G has a natural involution which is given by the map $\ast : g \mapsto g^{-1}$.

Definition

Let KG be the group algebra of a group G over a field K . If G is endowed with an involution \star , its linear extension to the group algebra KG is called a *K -linear involution* of KG . In particular, if $\star = \ast$ the induced involution is called *the classical involution*.

Canonical and K -linear involutions

Let G be a group endowed with an involution \star .

Let us consider the K -linear extension of \star to KG by setting

$$\left(\sum_{g \in G} a_g g \right)^\star := \sum_{g \in G} a_g g^\star.$$

This extension, which we denote again by \star , is an involution of KG which fixes the ground field K elementwise.

As well-known, any group G has a natural involution which is given by the map $\ast : g \mapsto g^{-1}$.

Definition

Let KG be the group algebra of a group G over a field K . If G is endowed with an involution \ast , its linear extension to the group algebra KG is called a *K -linear involution* of KG . In particular, if $\star = \ast$ the induced involution is called *the classical involution*.

Canonical and K -linear involutions

Let G be a group endowed with an involution \star .

Let us consider the K -linear extension of \star to KG by setting

$$\left(\sum_{g \in G} a_g g \right)^\star := \sum_{g \in G} a_g g^\star.$$

This extension, which we denote again by \star , is an involution of KG which fixes the ground field K elementwise.

As well-known, any group G has a natural involution which is given by the map $\ast : g \mapsto g^{-1}$.

Definition

Let KG be the group algebra of a group G over a field K . If G is endowed with an involution \star , its linear extension to the group algebra KG is called a *K -linear involution* of KG . In particular, if $\star = \ast$ the induced involution is called *the classical involution*.

Symmetric and skew-symmetric elements

Let KG be a group algebra endowed with a K -linear involution

\star .

Definition

Let KG be the group algebra of a group G over a field K endowed with a K -linear involution. The elements of KG^+ are called the *symmetric elements* of KG (with respect to \star) and those of KG^- are called the *skew-symmetric elements* of KG .

Symmetric and skew-symmetric elements

Let KG be a group algebra endowed with a K -linear involution \star . Let us set

$$KG^+ := \{x \mid x \in KG \quad x^\star = x\},$$

Definition

Let KG be the group algebra of a group G over a field K endowed with a K -linear involution. The elements of KG^+ are called the *symmetric elements* of KG (with respect to \star) and those of KG^- are called the *skew-symmetric elements* of KG .

Symmetric and skew-symmetric elements

Let KG be a group algebra endowed with a K -linear involution \star . Let us set

$$KG^+ := \{x \mid x \in KG \quad x^\star = x\},$$

$$KG^- := \{x \mid x \in KG \quad x^\star = -x\}.$$

Definition

Let KG be the group algebra of a group G over a field K endowed with a K -linear involution. The elements of KG^+ are called the *symmetric elements* of KG (with respect to \star) and those of KG^- are called the *skew-symmetric elements* of KG .

Symmetric and skew-symmetric elements

Let KG be a group algebra endowed with a K -linear involution \star . Let us set

$$KG^+ := \{x \mid x \in KG \quad x^\star = x\},$$

$$KG^- := \{x \mid x \in KG \quad x^\star = -x\}.$$

KG^- is a Lie subalgebra of KG , whereas KG^+ is a Jordan subalgebra of KG .

Definition

Let KG be the group algebra of a group G over a field K endowed with a K -linear involution. The elements of KG^+ are called the *symmetric elements* of KG (with respect to \star) and those of KG^- are called the *skew-symmetric elements* of KG .

Symmetric and skew-symmetric elements

Let KG be a group algebra endowed with a K -linear involution \star . Let us set

$$KG^+ := \{x \mid x \in KG \quad x^\star = x\},$$

$$KG^- := \{x \mid x \in KG \quad x^\star = -x\}.$$

KG^- is a Lie subalgebra of KG , whereas KG^+ is a Jordan subalgebra of KG .

Definition

Let KG be the group algebra of a group G over a field K endowed with a K -linear involution. The elements of KG^+ are called the *symmetric elements* of KG (with respect to \star) and those of KG^- are called the *skew-symmetric elements* of KG .

Symmetric and unitary units

Let us set

$$\mathcal{U}^+(KG) := \mathcal{U}(KG) \cap KG^+,$$

Definition

Let KG be the group algebra of a group G over a field K endowed with a K -linear involution. The elements of $\mathcal{U}^+(KG)$ are called the *symmetric units* of KG (with respect to \star) and those of $Un(KG)$ are called the *unitary units* of KG .

Symmetric and unitary units

Let us set

$$\mathcal{U}^+(KG) := \mathcal{U}(KG) \cap KG^+,$$

$$Un(KG) := \{x \mid x \in KG \quad xx^* = x^*x = 1\}.$$

Definition

Let KG be the group algebra of a group G over a field K endowed with a K -linear involution. The elements of $\mathcal{U}^+(KG)$ are called the *symmetric units* of KG (with respect to \star) and those of $Un(KG)$ are called the *unitary units* of KG .

Symmetric and unitary units

Let us set

$$\mathcal{U}^+(KG) := \mathcal{U}(KG) \cap KG^+,$$

$$Un(KG) := \{x \mid x \in KG \quad xx^* = x^*x = 1\}.$$

$Un(KG)$ is a subgroup of $\mathcal{U}(KG)$, whereas $\mathcal{U}^+(KG)$ is simply a subset of $\mathcal{U}(KG)$.

Definition

Let KG be the group algebra of a group G over a field K endowed with a K -linear involution. The elements of $\mathcal{U}^+(KG)$ are called the *symmetric units* of KG (with respect to \star) and those of $Un(KG)$ are called the *unitary units* of KG .

Symmetric and unitary units

Let us set

$$\mathcal{U}^+(KG) := \mathcal{U}(KG) \cap KG^+,$$

$$Un(KG) := \{x \mid x \in KG \quad xx^* = x^*x = 1\}.$$

$Un(KG)$ is a subgroup of $\mathcal{U}(KG)$, whereas $\mathcal{U}^+(KG)$ is simply a subset of $\mathcal{U}(KG)$.

Definition

Let KG be the group algebra of a group G over a field K endowed with a K -linear involution. The elements of $\mathcal{U}^+(KG)$ are called the *symmetric units* of KG (with respect to \star) and those of $Un(KG)$ are called the *unitary units* of KG .

Main question for symmetric and skew-symmetric elements

Main question for symmetric and skew-symmetric elements

First Question

To determine the extent to which the properties of the symmetric (or skew-symmetric) elements determine the properties of the whole group ring.

Main question for symmetric and skew-symmetric elements

First Question

To determine the extent to which the properties of the symmetric (or skew-symmetric) elements determine the properties of the whole group ring.

If A is an arbitrary algebra with involution, let us define in the same manner A^+ and A^- .

Theorem [Amitsur, 1968]

Let A be an algebra with involution. If A^+ or A^- satisfies a polynomial identity, then A satisfies a polynomial identity.

Lie properties for KG^+ and KG^-

Lie properties for KG^+ and KG^-

Lie properties for KG^+ and KG^-

Assume that KG is endowed with a K -linear involution. If KG^+ and/or KG^- satisfies a Lie identity, what can you say about KG ?

Lie nilpotency for KG^+ and KG^-

Theorem [Giambruno-Sehgal, 1993]

Let KG be the group algebra of a group G without 2-elements over a field K of characteristic $p \neq 2$ endowed with the classical involution. Then KG^+ or KG^- is Lie nilpotent if, and only if, KG is Lie nilpotent.

Lie nilpotency for KG^+ and KG^-

Theorem [Giambruno-Sehgal, 1993]

Let KG be the group algebra of a group G without 2-elements over a field K of characteristic $p \neq 2$ endowed with the classical involution. Then KG^+ or KG^- is Lie nilpotent if, and only if, KG is Lie nilpotent.

- The previous result **does not hold** without the assumption on the order of the elements of G .
- Lee (1999) completed the classification with regard to KG^+ by showing that the result is heavily effected by the presence of Q_8 in G .
- Giambruno-Sehgal (2007) completed the classification with regard to KG^- .

Theorem [Giambruno-Polcino Milies-Sehgal, 2009]

Let KG be the group algebra of a group G without 2-elements over a field K of characteristic $p \neq 2$ endowed with a K -linear involution. Then KG^+ is Lie nilpotent if, and only if, KG is Lie nilpotent.

Theorem [Giambruno-Polcino Milies-Sehgal, 2009]

Let KG be the group algebra of a group G without 2-elements over a field K of characteristic $p \neq 2$ endowed with a K -linear involution. Then KG^+ is Lie nilpotent if, and only if, KG is Lie nilpotent.

- Lee-Sehgal-Spinelli (2009) completed the classification for arbitrary G .
- Giambruno-Polcino Milies-Sehgal (2010) studied the question for KG^- , when G is a torsion group without 2-elements.

Lie n -Engel condition for KG^+ and KG^-

Theorem [Lee, 2000]

Let KG be the group algebra of a group G with no 2-elements over a field K of characteristic $p \neq 2$ endowed with the classical involution. Then KG^+ or KG^- is Lie n -Engel, for some n , if, and only if, KG is Lie m -Engel, for some m .

Lie n -Engel condition for KG^+ and KG^-

Theorem [Lee, 2000]

Let KG be the group algebra of a group G with no 2-elements over a field K of characteristic $p \neq 2$ endowed with the classical involution. Then KG^+ or KG^- is Lie n -Engel, for some n , if, and only if, KG is Lie m -Engel, for some m .

- Lee (2000) completed the classification with regard to KG^+ for arbitrary G .

Lie n -Engel condition for KG^+ and KG^-

Theorem [Lee, 2000]

Let KG be the group algebra of a group G with no 2-elements over a field K of characteristic $p \neq 2$ endowed with the classical involution. Then KG^+ or KG^- is Lie n -Engel, for some n , if, and only if, KG is Lie m -Engel, for some m .

- Lee (2000) completed the classification with regard to KG^+ for arbitrary G .

Theorem [Giambruno-Polcino Milies-Sehgal, 2009]

Let KG be the group algebra of a group G with no 2-elements over a field K of characteristic $p \neq 2$ endowed with a K -linear involution. Then KG^+ is Lie n -Engel, for some n , if, and only if, KG is Lie m -Engel, for some m .

Lie n -Engel condition for KG^+ and KG^-

Theorem [Lee, 2000]

Let KG be the group algebra of a group G with no 2-elements over a field K of characteristic $p \neq 2$ endowed with the classical involution. Then KG^+ or KG^- is Lie n -Engel, for some n , if, and only if, KG is Lie m -Engel, for some m .

- Lee (2000) completed the classification with regard to KG^+ for arbitrary G .

Theorem [Giambruno-Polcino Milies-Sehgal, 2009]

Let KG be the group algebra of a group G with no 2-elements over a field K of characteristic $p \neq 2$ endowed with a K -linear involution. Then KG^+ is Lie n -Engel, for some n , if, and only if, KG is Lie m -Engel, for some m .

- Lee-Sehgal-Spinelli (2009) completed the classification for

Lie solvability for KG^+ and KG^-

Remark

Let KG be the group algebra of a group G over a field K of characteristic $p \neq 2$. If KG^- is Lie solvable, then KG^+ is Lie solvable and

$$dl_L(KG^+) \leq dl_L(KG^-) + 1.$$

Lie solvability for KG^+ and KG^-

Remark

Let KG be the group algebra of a group G over a field K of characteristic $p \neq 2$. If KG^- is Lie solvable, then KG^+ is Lie solvable and

$$dl_L(KG^+) \leq dl_L(KG^-) + 1.$$

Lee, Sehgal and Spinelli studied the question for the classical involution.

Lie solvability for KG^+ and KG^-

Remark

Let KG be the group algebra of a group G over a field K of characteristic $p \neq 2$. If KG^- is Lie solvable, then KG^+ is Lie solvable and

$$dl_L(KG^+) \leq dl_L(KG^-) + 1.$$

Lee, Sehgal and Spinelli studied the question for the classical involution.

Theorem [Lee-Sehgal-S., 2009]

Let KG be the group algebra of a group G without 2-elements over a field K of characteristic $p \neq 2$ endowed with the classical involution. If P is finite, then KG^+ is Lie solvable if, and only if, KG is Lie solvable.

Theorem [Lee-Sehgal-S., 2009]

Let K be a field of characteristic $p > 2$, and let G be a group such that P contains an infinite subgroup of bounded exponent, and G contains no nontrivial elements of order dividing $p^2 - 1$. Then the following statements are equivalent:

- KG^- is Lie solvable;
- KG^+ is Lie solvable;
- KG is Lie solvable.

Theorem [Lee-Sehgal-S., 2009]

Let K be a field of characteristic $p > 2$, and let G be a group such that P contains an infinite subgroup of bounded exponent, and G contains no nontrivial elements of order dividing $p^2 - 1$. Then the following statements are equivalent:

- KG^- is Lie solvable;
- KG^+ is Lie solvable;
- KG is Lie solvable.

Theorem [Lee-Sehgal-S., 2009]

Let K be a field of characteristic $p > 2$ and G a group containing elements of infinite order, but no 2-elements. If KG^+ is Lie solvable, then KG is Lie solvable.

Main question for symmetric units

Main question for symmetric units

Second Question

To determine the extent to which the properties of the symmetric units determine the properties of the whole unit group of the group ring.

Main question for symmetric units

Second Question

To determine the extent to which the properties of the symmetric units determine the properties of the whole unit group of the group ring.

It is well-known that there is a strong connection between Lie properties satisfied by KG and the corresponding group identities satisfied by $\mathcal{U}(KG)$.

Main question for symmetric units

Second Question

To determine the extent to which the properties of the symmetric units determine the properties of the whole unit group of the group ring.

It is well-known that there is a strong connection between Lie properties satisfied by KG and the corresponding group identities satisfied by $\mathcal{U}(KG)$.

In this spirit is the following

Third Question

Do the Lie properties satisfied by the symmetric elements reflect the group identities satisfied by the symmetric units of the group ring?

When $\mathcal{U}^+(KG)$ is GI

When $\mathcal{U}^+(KG)$ is GI

Giambruno, Jespers, Sehgal, Valenti, Liu and Passman solved the Hartley's conjecture and classified when $\mathcal{U}(KG)$ is GI.

When $\mathcal{U}^+(KG)$ is GI

Giamb Bruno, Jespers, Sehgal, Valenti, Liu and Passman solved the Hartley's conjecture and classified when $\mathcal{U}(KG)$ is GI.

Theorem [Giamb Bruno-Sehgal-Valenti, 1998]

Let KG be the group algebra of a torsion group G over an infinite field K of characteristic $p \neq 2$.

- (a) If $p = 0$, $\mathcal{U}^+(KG)$ is GI if, and only if, G is either abelian or Hamiltonian 2-group.
- (b) If $p > 2$, $\mathcal{U}^+(KG)$ is GI if, and only if, KG is PI and either $Q_8 \not\subseteq G$ and G' is of bounded exponent p^k for some $k \geq 0$ or $Q_8 \subseteq G$ and
 - P is a normal subgroup of G and G/P is a Hamiltonian 2-group;
 - G is of bounded exponent $4p^s$ for some $s \geq 0$.

When $\mathcal{U}^+(KG)$ is GI

Giamb Bruno, Jespers, Sehgal, Valenti, Liu and Passman solved the Hartley's conjecture and classified when $\mathcal{U}(KG)$ is GI.

Theorem [Giamb Bruno-Sehgal-Valenti, 1998]

Let KG be the group algebra of a torsion group G over an infinite field K of characteristic $p \neq 2$.

- (a) If $p = 0$, $\mathcal{U}^+(KG)$ is GI if, and only if, G is either abelian or Hamiltonian 2-group.
- (b) If $p > 2$, $\mathcal{U}^+(KG)$ is GI if, and only if, KG is PI and either $Q_8 \not\subseteq G$ and G' is of bounded exponent p^k for some $k \geq 0$ or $Q_8 \subseteq G$ and
 - P is a normal subgroup of G and G/P is a Hamiltonian 2-group;
 - G is of bounded exponent $4p^s$ for some $s \geq 0$.

- Sehgal-Valenti (2006) studied the non-torsion case.

Special group identities

After the result by Giambruno, Sehgal and Valenti, it was of interest to consider when $\mathcal{U}^+(KG)$ satisfies special group identities.

Special group identities

After the result by Giambruno, Sehgal and Valenti, it was of interest to consider when $\mathcal{U}^+(KG)$ satisfies special group identities.

Theorem [Lee, 2003]

Let KG be the group algebra of a torsion group G over a field K of characteristic $p \neq 2$ endowed with the classical involution.

$$\mathcal{U}^+(KG) \text{ is nilpotent} \iff KG^+ \text{ is Lie nilpotent.}$$

Special group identities

After the result by Giambruno, Sehgal and Valenti, it was of interest to consider when $\mathcal{U}^+(KG)$ satisfies special group identities.

Theorem [Lee, 2003]

Let KG be the group algebra of a torsion group G over a field K of characteristic $p \neq 2$ endowed with the classical involution.

$$\mathcal{U}^+(KG) \text{ is nilpotent} \iff KG^+ \text{ is Lie nilpotent.}$$

- Lee-Polcino Milies-Sehgal (2007) studied the non-torsion case.

Other group identities

Theorem [Lee-Spinelli, 2010]

Let KG be the group algebra of a torsion group G over an infinite field K of characteristic $p \neq 2$ endowed with the classical involution.

$$\langle \mathcal{U}^+(KG) \rangle \text{ is Engel} \iff KG^+ \text{ is Lie Engel.}$$

Other group identities

Theorem [Lee-Spinelli, 2010]

Let KG be the group algebra of a torsion group G over an infinite field K of characteristic $p \neq 2$ endowed with the classical involution.

$$\langle \mathcal{U}^+(KG) \rangle \text{ is Engel} \iff KG^+ \text{ is Lie Engel.}$$

Theorem [Lee-Spinelli, 2009]

Let KG be the group algebra of a torsion group G over an infinite field K of characteristic $p \neq 2$ endowed with the classical involution. If P is infinite and G does not contain elements whose order divides $p^2 - 1$,

$$\mathcal{U}^+(KG) \text{ is solvable} \iff KG^+ \text{ is Lie solvable.}$$

When $\mathcal{U}^+(KG)$ is GI

When $\mathcal{U}^+(KG)$ is GI

After a paper by Dooms and Ruiz Marin (2007), the key result is the following

When $\mathcal{U}^+(KG)$ is GI

After a paper by Dooms and Ruiz Marin (2007), the key result is the following

Theorem [Giambruno-Polcino Milies-Sehgal, 2009]

Let KG be the group algebra of a torsion group G over an infinite field K of characteristic $p \neq 2$ endowed with a K -linear involution. Then $\mathcal{U}^+(KG)$ is GI if, and only if,

- (a) KG is semiprime and G is either abelian or an SLC -group, or
- (b) KG is not semiprime, P is a normal subgroup of G , G has a p -abelian normal subgroup of finite index and either
 - G' is a p -group of bounded exponent or
 - G/P is an SLC -group and G contains a normal $*$ -invariant p -subgroup B of bounded exponent such that P/B is central in G/P and the induced involution acts as the identity on P/B .

SLC-groups

SLC-groups

A group G is called an *LC-group* (that is, it has the “*lack of commutativity*” property) if it is not abelian, but, whenever $g, h \in G$ and $gh = hg$, then at least one of $\{g, h, gh\}$ is central.

SLC-groups

A group G is called an *LC-group* (that is, it has the “*lack of commutativity*” property) if it is not abelian, but, whenever $g, h \in G$ and $gh = hg$, then at least one of $\{g, h, gh\}$ is central. These groups were introduced by *E. Goodaire*.

SLC-groups

A group G is called an *LC-group* (that is, it has the “*lack of commutativity*” property) if it is not abelian, but, whenever $g, h \in G$ and $gh = hg$, then at least one of $\{g, h, gh\}$ is central. These groups were introduced by *E. Goodaire*.

A group G is an *LC-group* with a unique nonidentity commutator (which must, obviously, have order 2) if, and only if, $G/\zeta(G) \simeq C_2 \times C_2$.

SLC-groups

A group G is called an *LC-group* (that is, it has the “*lack of commutativity*” property) if it is not abelian, but, whenever $g, h \in G$ and $gh = hg$, then at least one of $\{g, h, gh\}$ is central. These groups were introduced by *E. Goodaire*.

A group G is an *LC-group* with a unique nonidentity commutator (which must, obviously, have order 2) if, and only if, $G/\zeta(G) \simeq C_2 \times C_2$.

Definition

A group G endowed with an involution \star is said to be a *special LC-group*, or *SLC-group*, if it is an *LC-group*, it has a unique nonidentity commutator z and, for all $g \in G$, we have $g^\star = g$ if $g \in \zeta(G)$ and, otherwise, $g^\star = zg$.

Theorem [Jespers-Ruiz Marin, 2006]

Let R be a commutative ring of characteristic different from 2, and G a nonabelian group endowed with an involution \star . Then RG^+ is commutative if, and only if, G is an SLC-group.

When $\mathcal{U}^+(KG)$ is nilpotent

Assume that $\mathcal{U}^+(KG)$ is nilpotent.

When $\mathcal{U}^+(KG)$ is nilpotent

Assume that $\mathcal{U}^+(KG)$ is nilpotent.

- Assume that KG is not semiprime, otherwise we are done by [GPMS] and [JRM].

When $\mathcal{U}^+(KG)$ is nilpotent

Assume that $\mathcal{U}^+(KG)$ is nilpotent.

- Assume that KG is not semiprime, otherwise we are done by [GPMS] and [JRM].
- By [GPMS] we know that $P \trianglelefteq G$.

When $U^+(KG)$ is nilpotent

Assume that $U^+(KG)$ is nilpotent.

- Assume that KG is not semiprime, otherwise we are done by [GPMS] and [JRM].
- By [GPMS] we know that $P \trianglelefteq G$.
- Let $N \trianglelefteq G$ and \star -invariant. If $U^+(KG)$ satisfies w , then $U^+(K(G/N))$ satisfies w .

When $\mathcal{U}^+(KG)$ is nilpotent

Assume that $\mathcal{U}^+(KG)$ is nilpotent.

- Assume that KG is not semiprime, otherwise we are done by [GPMS] and [JRM].
- By [GPMS] we know that $P \trianglelefteq G$.
- Let $N \trianglelefteq G$ and \star -invariant. If $\mathcal{U}^+(KG)$ satisfies w , then $\mathcal{U}^+(K(G/N))$ satisfies w .
- $\mathcal{U}^+(K(G/P))$ is nilpotent and $K(G/P)$ is semiprime.

When $\mathcal{U}^+(KG)$ is nilpotent

Assume that $\mathcal{U}^+(KG)$ is nilpotent.

- Assume that KG is not semiprime, otherwise we are done by [GPMS] and [JRM].
- By [GPMS] we know that $P \trianglelefteq G$.
- Let $N \trianglelefteq G$ and \star -invariant. If $\mathcal{U}^+(KG)$ satisfies w , then $\mathcal{U}^+(K(G/N))$ satisfies w .
- $\mathcal{U}^+(K(G/P))$ is nilpotent and $K(G/P)$ is semiprime.
- By [GPMS] G/P is abelian or G/P is an *SLC*-group.

G finite

G finite

By [GPMS] G is locally finite. Hence it is relevant to study the case in which G is finite.

Lemma

Let G be a finite group. If $\mathcal{U}^+(KG)$ is nilpotent, then G is nilpotent and G/P is either abelian or an *SLC*-group.

G/P abelian

G/P abelian

Theorem [Lee-Sehgal-S., 2010]

Let G/P be abelian. If $\mathcal{U}^+(KG)$ is nilpotent, then G is nilpotent and p -abelian (hence, $\mathcal{U}(KG)$ is nilpotent).

G/P is an *SLC*-group

G/P is an SLC -group

Theorem [Lee-Sehgal-S., 2010]

Let G/P be an SLC -group. Then $\mathcal{U}^+(KG)$ is nilpotent if, and only if, G is nilpotent and G has a finite normal $*$ -invariant p -subgroup N such that G/N is an SLC -group.

Main Theorem

Main Theorem

Theorem [Lee-Sehgal-S., 2010]

Let K be an infinite field of characteristic $p > 2$ and G a torsion group having an involution $*$, and let KG have the induced involution. Suppose that $U(KG)$ is not nilpotent. Then $U^+(KG)$ is nilpotent if, and only if, G is nilpotent and G has a finite normal $*$ -invariant p -subgroup N such that G/N is an *SLC*-group.

According to the result by Passi-Passman-Sehgal (1973) and Khripta (1972) KG is Lie nilpotent if, and only if, $\mathcal{U}(KG)$ is nilpotent.

According to the result by Passi-Passman-Sehgal (1973) and Khripta (1972) KG is Lie nilpotent if, and only if, $\mathcal{U}(KG)$ is nilpotent.

By using the results of [LSS1], one has

Theorem [Lee-Sehgal-S., 2010]

Let K be an infinite field of characteristic $p \neq 2$ and G a torsion group having an involution $*$, and let KG have the induced involution. Then $\mathcal{U}^+(KG)$ is nilpotent if, and only if, KG^+ is Lie nilpotent.