

On Duo Group Rings

Yuanlin Li

Brock University



Contents

- 1 Introduction and Preliminaries
- 2 Group Algebras KG
 - Characteristic 0
 - Characteristic 2
 - General Case
- 3 Group rings over integral domains
 - When is RQ_8 duo?
 - When is RG duo?
- 4 References

Introduction and Preliminaries

● Some Notations

1. R – Associative ring with 1.
2. kG – Group ring over a commutative ring k .
3. $Q_8 = \langle a, b \mid a^4 = 1, a^2 = b^2, bab^{-1} = a^{-1} \rangle$.

○ Definition 1.1

Let R be an associative ring with identity. Call R left (right) duo if every left (right) ideal is an ideal, and call R duo if it is both left and right duo. Define R to be reversible if $\alpha\beta = 0$ implies $\beta\alpha = 0$, and symmetric if $\alpha\beta\gamma = 0$ implies $\alpha\gamma\beta = 0$ for all $\alpha, \beta, \gamma \in R$. Finally, say that R has the “SI” property if $\alpha\beta = 0$ implies $\alpha R\beta = 0$ for all $\alpha, \beta \in R$.



Introduction and Preliminaries

● Some Notations

1. R – Associative ring with 1.
2. kG – Group ring over a commutative ring k .
3. $Q_8 = \langle a, b \mid a^4 = 1, a^2 = b^2, bab^{-1} = a^{-1} \rangle$.

● Definition 1.1

Let R be an associative ring with identity. Call R left (right) duo if every left (right) ideal is an ideal, and call R duo if it is both left and right duo. Define R to be reversible if $\alpha\beta = 0$ implies $\beta\alpha = 0$, and symmetric if $\alpha\beta\gamma = 0$ implies $\alpha\gamma\beta = 0$ for all $\alpha, \beta, \gamma \in R$. Finally, say that R has the “SI” property if $\alpha\beta = 0$ implies $\alpha R\beta = 0$ for all $\alpha, \beta \in R$.

Introduction and Preliminaries

- **Some Notations**

1. R – Associative ring with 1.
2. kG – Group ring over a commutative ring k .
3. $Q_8 = \langle a, b \mid a^4 = 1, a^2 = b^2, bab^{-1} = a^{-1} \rangle$.

- **Definition 1.1**

Let R be an associative ring with identity. Call R left (right) duo if every left (right) ideal is an ideal, and call R duo if it is both left and right duo. Define R to be reversible if $\alpha\beta = 0$ implies $\beta\alpha = 0$, and symmetric if $\alpha\beta\gamma = 0$ implies $\alpha\gamma\beta = 0$ for all $\alpha, \beta, \gamma \in R$. Finally, say that R has the “SI” property if $\alpha\beta = 0$ implies $\alpha R \beta = 0$ for all $\alpha, \beta \in R$.

Introduction and Preliminaries

- **Some Notations**

1. R – Associative ring with 1.
2. kG – Group ring over a commutative ring k .
3. $Q_8 = \langle a, b \mid a^4 = 1, a^2 = b^2, bab^{-1} = a^{-1} \rangle$.

- **Definition 1.1**

Let R be an associative ring with identity. Call R left (right) duo if every left (right) ideal is an ideal, and call R duo if it is both left and right duo. Define R to be reversible if $\alpha\beta = 0$ implies $\beta\alpha = 0$, and symmetric if $\alpha\beta\gamma = 0$ implies $\alpha\gamma\beta = 0$ for all $\alpha, \beta, \gamma \in R$. Finally, say that R has the “SI” property if $\alpha\beta = 0$ implies $\alpha R\beta = 0$ for all $\alpha, \beta \in R$.

Introduction and Preliminaries

- **Some Notations**

1. R – Associative ring with 1.
2. kG – Group ring over a commutative ring k .
3. $Q_8 = \langle a, b \mid a^4 = 1, a^2 = b^2, bab^{-1} = a^{-1} \rangle$.

- **Definition 1.1**

Let R be an associative ring with identity. Call R left (right) duo if every left (right) ideal is an ideal, and call R duo if it is both left and right duo. Define R to be reversible if $\alpha\beta = 0$ implies $\beta\alpha = 0$, and symmetric if $\alpha\beta\gamma = 0$ implies $\alpha\gamma\beta = 0$ for all $\alpha, \beta, \gamma \in R$. Finally, say that R has the “SI” property if $\alpha\beta = 0$ implies $\alpha R\beta = 0$ for all $\alpha, \beta \in R$.

Introduction and Preliminaries

- **Remark** Let k be a commutative ring with identity and G be any group. Using the standard involution $*$ on the group ring kG , defined by $(\sum a_i g_i)^* = \sum a_i g_i^{-1}$ for all $a_i \in k$ and $g_i \in G$, we see that kG is left duo if and only if it is right duo.
- Marks [5] has clarified the relationships among duo, reversible and symmetric rings. Moreover, he proved the following result [5, Proposition 6].
- **Proposition 1.2** Let k be a commutative ring with identity, and let G be a finite group. Then the group ring kG is reversible if and only if kG has the “SI” property.

Introduction and Preliminaries

- **Remark** Let k be a commutative ring with identity and G be any group. Using the standard involution $*$ on the group ring kG , defined by $(\sum a_i g_i)^* = \sum a_i g_i^{-1}$ for all $a_i \in k$ and $g_i \in G$, we see that kG is left duo if and only if it is right duo.
- Marks [5] has clarified the relationships among duo, reversible and symmetric rings. Moreover, he proved the following result [5, Proposition 6].
- **Proposition 1.2** Let k be a commutative ring with identity, and let G be a finite group. Then the group ring kG is reversible if and only if kG has the “SI” property.

Introduction and Preliminaries

- **Remark** Let k be a commutative ring with identity and G be any group. Using the standard involution $*$ on the group ring kG , defined by $(\sum a_i g_i)^* = \sum a_i g_i^{-1}$ for all $a_i \in k$ and $g_i \in G$, we see that kG is left duo if and only if it is right duo.
- **Marks [5]** has clarified the relationships among duo, reversible and symmetric rings. Moreover, he proved the following result [5, Proposition 6].
- **Proposition 1.2** Let k be a commutative ring with identity, and let G be a finite group. Then the group ring kG is reversible if and only if kG has the “SI” property.

Introduction and Preliminaries

- **Remark** Let k be a commutative ring with identity and G be any group. Using the standard involution $*$ on the group ring kG , defined by $(\sum a_i g_i)^* = \sum a_i g_i^{-1}$ for all $a_i \in k$ and $g_i \in G$, we see that kG is left duo if and only if it is right duo.
- Marks [5] has clarified the relationships among duo, reversible and symmetric rings. Moreover, he proved the following result [5, Proposition 6].
- **Proposition 1.2** Let k be a commutative ring with identity, and let G be a finite group. Then the group ring kG is reversible if and only if kG has the “SI” property.

Introduction and Preliminaries

- **Remark** Let k be a commutative ring with identity and G be any group. Using the standard involution $*$ on the group ring kG , defined by $(\sum a_i g_i)^* = \sum a_i g_i^{-1}$ for all $a_i \in k$ and $g_i \in G$, we see that kG is left duo if and only if it is right duo.
- Marks [5] has clarified the relationships among duo, reversible and symmetric rings. Moreover, he proved the following result [5, Proposition 6].
- **Proposition 1.2** Let k be a commutative ring with identity, and let G be a finite group. Then the group ring kG is reversible if and only if kG has the “SI” property.

Introduction and Preliminaries

- **Remark** Let k be a commutative ring with identity and G be any group. Using the standard involution $*$ on the group ring kG , defined by $(\sum a_i g_i)^* = \sum a_i g_i^{-1}$ for all $a_i \in k$ and $g_i \in G$, we see that kG is left duo if and only if it is right duo.
- Marks [5] has clarified the relationships among duo, reversible and symmetric rings. Moreover, he proved the following result [5, Proposition 6].
- **Proposition 1.2** Let k be a commutative ring with identity, and let G be a finite group. Then the group ring kG is reversible if and only if kG has the “SI” property.

Introduction and Preliminaries

- We note that this result remains valid for an arbitrary group G .
- We also note that the “SI” property is simply the statement that left annihilators and right annihilators are ideals, hence it is obvious that duo rings have the “SI” property. It now follows from Proposition 1.2 that if kG is a duo ring, then it is reversible.
- However, the converse is not true, as the following example shows.

Introduction and Preliminaries

- We note that this result remains valid for an arbitrary group G .
- We also note that the “SI” property is simply the statement that left annihilators and right annihilators are ideals, hence it is obvious that duo rings have the “SI” property. It now follows from Proposition 1.2 that if kG is a duo ring, then it is reversible.
- However, the converse is not true, as the fol-



Introduction and Preliminaries

- We note that this result remains valid for an arbitrary group G .
- We also note that the “SI” property is simply the statement that left annihilators and right annihilators are ideals, hence it is obvious that duo rings have the “SI” property. It now follows from Proposition 1.2 that if kG is a duo ring, then it is reversible.
- However, the converse is not true, as the fol-

Introduction and Preliminaries

- We note that this result remains valid for an arbitrary group G .
- We also note that the “SI” property is simply the statement that left annihilators and right annihilators are ideals, hence it is obvious that duo rings have the “SI” property. It now follows from Proposition 1.2 that if kG is a duo ring, then it is reversible.
- **However, the converse is not true, as the following example shows.**

Introduction and Preliminaries

- We note that this result remains valid for an arbitrary group G .
- We also note that the “SI” property is simply the statement that left annihilators and right annihilators are ideals, hence it is obvious that duo rings have the “SI” property. It now follows from Proposition 1.2 that if kG is a duo ring, then it is reversible.
- However, the converse is not true, as the following example shows.

Introduction and Preliminaries

- We note that this result remains valid for an arbitrary group G .
- We also note that the “SI” property is simply the statement that left annihilators and right annihilators are ideals, hence it is obvious that duo rings have the “SI” property. It now follows from Proposition 1.2 that if kG is a duo ring, then it is reversible.
- However, the converse is not true, as the following example shows.

Introduction and Preliminaries

- **Example 1.3** Let $Q_8 = \langle a, b | a^4 = 1, a^2 = b^2, a^b = a^{-1} \rangle$ be the quaternion group of order 8. The integral group ring $\mathbb{Z}Q_8$ is a reversible ring, but not a duo ring.
- **Proof.** It follows from Theorem 3.1 in [3] that the rational group algebra $\mathbb{Q}Q_8$ is reversible. As a subring of $\mathbb{Q}Q_8$, $\mathbb{Z}Q_8$ is clearly reversible.
- $R = \mathbb{Z}Q_8$ is not a duo ring since the left ideal $R(a + 2b)$ generated by $a + 2b$ is not a right ideal.

Introduction and Preliminaries

- **Example 1.3** Let $Q_8 = \langle a, b | a^4 = 1, a^2 = b^2, a^b = a^{-1} \rangle$ be the quaternion group of order 8. The integral group ring $\mathbb{Z}Q_8$ is a reversible ring, but not a duo ring.
- **Proof.** It follows from Theorem 3.1 in [3] that the rational group algebra $\mathbb{Q}Q_8$ is reversible. As a subring of $\mathbb{Q}Q_8$, $\mathbb{Z}Q_8$ is clearly reversible.
- $R = \mathbb{Z}Q_8$ is not a duo ring since the left ideal $R(a + 2b)$ generated by $a + 2b$ is not a right



Introduction and Preliminaries

- **Example 1.3** Let $Q_8 = \langle a, b \mid a^4 = 1, a^2 = b^2, a^b = a^{-1} \rangle$ be the quaternion group of order 8. The integral group ring $\mathbb{Z}Q_8$ is a reversible ring, but not a duo ring.
- **Proof.** It follows from Theorem 3.1 in [3] that the rational group algebra $\mathbb{Q}Q_8$ is reversible. As a subring of $\mathbb{Q}Q_8$, $\mathbb{Z}Q_8$ is clearly reversible.
- $R = \mathbb{Z}Q_8$ is not a duo ring since the left ideal $R(a + 2b)$ generated by $a + 2b$ is not a right ideal.

Introduction and Preliminaries

- **Example 1.3** Let $Q_8 = \langle a, b \mid a^4 = 1, a^2 = b^2, a^b = a^{-1} \rangle$ be the quaternion group of order 8. The integral group ring $\mathbb{Z}Q_8$ is a reversible ring, but not a duo ring.
- **Proof.** It follows from Theorem 3.1 in [3] that the rational group algebra $\mathbb{Q}Q_8$ is reversible. As a subring of $\mathbb{Q}Q_8$, $\mathbb{Z}Q_8$ is clearly reversible.
- $R = \mathbb{Z}Q_8$ is not a duo ring since the left ideal $R(a + 2b)$ generated by $a + 2b$ is not a right ideal.

Introduction and Preliminaries

- **Example 1.3** Let $Q_8 = \langle a, b \mid a^4 = 1, a^2 = b^2, a^b = a^{-1} \rangle$ be the quaternion group of order 8. The integral group ring $\mathbb{Z}Q_8$ is a reversible ring, but not a duo ring.
- **Proof.** It follows from Theorem 3.1 in [3] that the rational group algebra $\mathbb{Q}Q_8$ is reversible. As a subring of $\mathbb{Q}Q_8$, $\mathbb{Z}Q_8$ is clearly reversible.
- $R = \mathbb{Z}Q_8$ is not a duo ring since the left ideal $R(a + 2b)$ generated by $a + 2b$ is not a right ideal.

Introduction and Preliminaries

- **Example 1.3** Let $Q_8 = \langle a, b \mid a^4 = 1, a^2 = b^2, a^b = a^{-1} \rangle$ be the quaternion group of order 8. The integral group ring $\mathbb{Z}Q_8$ is a reversible ring, but not a duo ring.
- **Proof.** It follows from Theorem 3.1 in [3] that the rational group algebra $\mathbb{Q}Q_8$ is reversible. As a subring of $\mathbb{Q}Q_8$, $\mathbb{Z}Q_8$ is clearly reversible.
- $R = \mathbb{Z}Q_8$ is not a duo ring since the left ideal $R(a + 2b)$ generated by $a + 2b$ is not a right ideal.

Introduction and Preliminaries

- In this talk, we shall study when the group rings RG of torsion groups over integral domains are duo.
- If $R = K$ is a field, then that KG is a duo ring if and only if KG is reversible.
- If G is a non-abelian torsion group and R is an ID, then RG is a duo ring if and only if R is a field and RG is reversible for the following cases: (1) $\text{char } R \neq 0$, and (2) $\text{char } R = 0$, and $S \subseteq R \subseteq K$, where S is a ring of algebraic integers.



Introduction and Preliminaries

- In this talk, we shall study when the group rings RG of torsion groups over integral domains are duo.
- If $R = K$ is a field, then that KG is a duo ring if and only if KG is reversible.
- If G is a non-abelian torsion group and R is an ID, then RG is a duo ring if and only if R is a field and RG is reversible for the following cases: (1) $\text{char } R \neq 0$, and (2) $\text{char } R = 0$, and $S \subseteq R \subseteq K$, where S is a ring of algebraic

Introduction and Preliminaries

- In this talk, we shall study when the group rings RG of torsion groups over integral domains are duo.
- If $R = K$ is a field, then that KG is a duo ring if and only if KG is reversible.
- If G is a non-abelian torsion group and R is an ID, then RG is a duo ring if and only if R is a field and RG is reversible for the following cases: (1) $\text{char } R \neq 0$, and (2) $\text{char } R = 0$, and $S \subseteq R \subseteq K_S$, where S is a ring of algebraic integers and K_S is its quotient field.

Introduction and Preliminaries

- In this talk, we shall study when the group rings RG of torsion groups over integral domains are duo.
- If $R = K$ is a field, then that KG is a duo ring if and only if KG is reversible.
- If G is a non-abelian torsion group and R is an ID, then RG is a duo ring if and only if R is a field and RG is reversible for the following cases: (1) $\text{char } R \neq 0$, and (2) $\text{char } R = 0$, and $S \subseteq R \subseteq K_S$, where S is a ring of algebraic integers and K_S is its quotient field.

Introduction and Preliminaries

- In this talk, we shall study when the group rings RG of torsion groups over integral domains are duo.
- If $R = K$ is a field, then that KG is a duo ring if and only if KG is reversible.
- If G is a non-abelian torsion group and R is an ID, then RG is a duo ring if and only if R is a field and RG is reversible for the following cases: (1) $\text{char } R \neq 0$, and (2) $\text{char } R = 0$, and $S \subseteq R \subseteq K_S$, where S is a ring of algebraic integers and K_S is its quotient field.

Introduction and Preliminaries

- In this talk, we shall study when the group rings RG of torsion groups over integral domains are duo.
- If $R = K$ is a field, then that KG is a duo ring if and only if KG is reversible.
- If G is a non-abelian torsion group and R is an ID, then RG is a duo ring if and only if R is a field and RG is reversible for the following cases: (1) $\text{char } R \neq 0$, and (2) $\text{char } R = 0$, and $S \subseteq R \subseteq K_S$, where S is a ring of algebraic integers and K_S is its quotient field.

Introduction and Preliminaries

- It was shown in [3, 4] that if RG of a torsion group G over a commutative ring R is reversible, then either G is an abelian group, or G is a Hamiltonian group. In addition, if $R = K$ is a field then the characteristic of K is 0 or 2.
- Next assume that $\text{char}(R) = 0$ or 2 and G is a Hamiltonian group, i.e. $G = Q_8 \times E_2 \times E'_2$.
- We first deal with the group algebra KG case where K is a field. The general case RG will be handled in the last section.

Introduction and Preliminaries

- It was shown in [3, 4] that if RG of a torsion group G over a commutative ring R is reversible, then either G is an abelian group, or G is a Hamiltonian group. In addition, if $R = K$ is a field then the characteristic of K is 0 or 2.
- Next assume that $\text{char}(R) = 0$ or 2 and G is a Hamiltonian group, i.e. $G = Q_8 \times E_2 \times E'_2$.
- We first deal with the group algebra KG case where K is a field. The general case RG will

Introduction and Preliminaries

- It was shown in [3, 4] that if RG of a torsion group G over a commutative ring R is reversible, then either G is an abelian group, or G is a Hamiltonian group. In addition, if $R = K$ is a field then the characteristic of K is 0 or 2.
- Next assume that $\text{char}(R) = 0$ or 2 and G is a Hamiltonian group, i.e. $G = Q_8 \times E_2 \times E'_2$.
- We first deal with the group algebra KG case where K is a field. The general case RG will

Introduction and Preliminaries

- It was shown in [3, 4] that if RG of a torsion group G over a commutative ring R is reversible, then either G is an abelian group, or G is a Hamiltonian group. In addition, if $R = K$ is a field then the characteristic of K is 0 or 2.
- Next assume that $\text{char}(R) = 0$ or 2 and G is a Hamiltonian group, i.e. $G = Q_8 \times E_2 \times E'_2$.
- We first deal with the group algebra KG case where K is a field. The general case RG will be handled in the last section.

Introduction and Preliminaries

- It was shown in [3, 4] that if RG of a torsion group G over a commutative ring R is reversible, then either G is an abelian group, or G is a Hamiltonian group. In addition, if $R = K$ is a field then the characteristic of K is 0 or 2.
- Next assume that $\text{char}(R) = 0$ or 2 and G is a Hamiltonian group, i.e. $G = Q_8 \times E_2 \times E'_2$.
- We first deal with the group algebra KG case where K is a field. The general case RG will be handled in the last section.

Introduction and Preliminaries

- It was shown in [3, 4] that if RG of a torsion group G over a commutative ring R is reversible, then either G is an abelian group, or G is a Hamiltonian group. In addition, if $R = K$ is a field then the characteristic of K is 0 or 2.
- Next assume that $\text{char}(R) = 0$ or 2 and G is a Hamiltonian group, i.e. $G = Q_8 \times E_2 \times E'_2$.
- We first deal with the group algebra KG case where K is a field. The general case RG will be handled in the last section.

Group Algebras KG

- 1 Let $G = Q_8$ and K be a field of characteristic 0. The following result can be derived from some known results in group rings along with the Wedderburn decomposition of KQ_8 .
- 2 **Theorem 2.1** The following statements are equivalent:
 - (1) $R = KQ_8$ is duo.
 - (2) $R = KQ_8$ is reversible.
 - (3) The equation $1 + x^2 + y^2 = 0$ has no solutions in K .



Group Algebras KG

- 1 Let $G = Q_8$ and K be a field of characteristic 0. The following result can be derived from some known results in group rings along with the Wedderburn decomposition of KQ_8 .
- 2 **Theorem 2.1** The following statements are equivalent:
 - (1) $R = KQ_8$ is duo.
 - (2) $R = KQ_8$ is reversible.
 - (3) The equation $1 + x^2 + y^2 = 0$ has no solutions in K .

Group Algebras KG

- 1 Let $G = Q_8$ and K be a field of characteristic 0. The following result can be derived from some known results in group rings along with the Wedderburn decomposition of KQ_8 .
- 2 **Theorem 2.1** The following statements are equivalent:
 - (1) $R = KQ_8$ is duo.
 - (2) $R = KQ_8$ is reversible.
 - (3) The equation $1 + x^2 + y^2 = 0$ has no solutions in K .

Group Algebras KG

- We now characterize when KQ_8 is duo, where K is a field with characteristic 2.
- **Theorem 2.2** The following statements are equivalent:
 - (1) KQ_8 is duo.
 - (2) KQ_8 is reversible.
 - (3) The equation $1 + x + x^2 = 0$ has no solutions in K .

Group Algebras KG

- We now characterize when KQ_8 is duo, where K is a field with characteristic 2.
- **Theorem 2.2** The following statements are equivalent:
 - (1) KQ_8 is duo.
 - (2) KQ_8 is reversible.
 - (3) The equation $1 + x + x^2 = 0$ has no solutions in K .

Group Algebras KG

- **Theorem 2.5** Let K be a field and let G be a torsion group. Then KG is a duo ring if and only if one of the following conditions holds.
 - (1) G is Abelian.
 - (2) $G = Q_8 \times E_2 \times E'_2$ is Hamiltonian, the characteristic of K is 0 and the equation $1 + x^2 + y^2 = 0$ has no solutions in any cyclotomic field $K(\xi_d)$ for any odd d which is an order of an element of E'_2 .
 - (3) $G = Q_8 \times E'_2$, the characteristic of K is 2 and the equation $1 + x + x^2 = 0$ has no solutions in any cyclotomic field $K(\xi_d)$ for any odd d which is an order of an element of E'_2 .

Group Algebras KG

- **Theorem 2.5** Let K be a field and let G be a torsion group. Then KG is a duo ring if and only if one of the following conditions holds.
 - (1) G is Abelian.
 - (2) $G = Q_8 \times E_2 \times E'_2$ is Hamiltonian, the characteristic of K is 0 and the equation $1 + x^2 + y^2 = 0$ has no solutions in any cyclotomic field $K(\xi_d)$ for any odd d which is an order of an element of E'_2 .
 - (3) $G = Q_8 \times E'_2$, the characteristic of K is 2 and the equation $1 + x + x^2 = 0$ has no solutions in any cyclotomic field $K(\xi_d)$ for any odd d which is an order of an element of E'_2 .

Group Algebras KG

① **Corollary 2.6** KG is duo if and only if KG is reversible.

Group Algebras KG

① **Corollary 2.6** KG is duo if and only if KG is reversible.

When is RQ_8 duo?

Group rings over integral domains

- ① We remark that if G is a non-abelian torsion group and R is a commutative ring with identity for which RG is duo, then RQ_8 is also duo. The question of when RG is duo essentially reduces to that of when RQ_8 is duo, where R is an integral domain.
- As mentioned early, the integral group ring $\mathbb{Z}Q_8$ is a reversible ring, but not a duo ring [1, Example 1.1] while $\mathbb{Q}Q_8$ is a duo ring. A natural question which arises is as follows:
- **Question 3.1** Is there any ring R between \mathbb{Z} and \mathbb{Q} (in addition to \mathbb{Q} the field of all rational numbers), such that RQ_8 is duo.



When is RQ_8 duo?

Group rings over integral domains

- ① We remark that if G is a non-abelian torsion group and R is a commutative ring with identity for which RG is duo, then RQ_8 is also duo. The question of when RG is duo essentially reduces to that of when RQ_8 is duo, where R is an integral domain.
- As mentioned early, the integral group ring $\mathbb{Z}Q_8$ is a reversible ring, but not a duo ring [1, Example 1.1] while $\mathbb{Q}Q_8$ is a duo ring. A natural question which arises is as follows:
- **Question 3.1** Is there any ring R between \mathbb{Z} and \mathbb{Q} (in addition to \mathbb{Q} the field of all rational numbers), such that RQ_8 is duo.



When is RQ_8 duo?

Group rings over integral domains

- 1 We remark that if G is a non-abelian torsion group and R is a commutative ring with identity for which RG is duo, then RQ_8 is also duo. The question of when RG is duo essentially reduces to that of when RQ_8 is duo, where R is an integral domain.
- As mentioned early, the integral group ring $\mathbb{Z}Q_8$ is a reversible ring, but not a duo ring [1, Example 1.1] while $\mathbb{Q}Q_8$ is a duo ring. A natural question which arises is as follows:
- Question 3.1** Is there any ring R between \mathbb{Z} and \mathbb{Q} (in addition to \mathbb{Q} the field of all rational numbers), such that RQ_8 is duo.



When is RQ_8 duo?

Group rings over integral domains

- 1 We remark that if G is a non-abelian torsion group and R is a commutative ring with identity for which RG is duo, then RQ_8 is also duo. The question of when RG is duo essentially reduces to that of when RQ_8 is duo, where R is an integral domain.
- 2 As mentioned early, the integral group ring $\mathbb{Z}Q_8$ is a reversible ring, but not a duo ring [1, Example 1.1] while $\mathbb{Q}Q_8$ is a duo ring. A natural question which arises is as follows:
 - Question 3.1 Is there any ring R between \mathbb{Z} and \mathbb{Q} (in addition to \mathbb{Q} the field of all rational numbers), such



When is RQ_8 duo?

Group rings over integral domains

- 1 We remark that if G is a non-abelian torsion group and R is a commutative ring with identity for which RG is duo, then RQ_8 is also duo. The question of when RG is duo essentially reduces to that of when RQ_8 is duo, where R is an integral domain.
- 2 As mentioned early, the integral group ring $\mathbb{Z}Q_8$ is a reversible ring, but not a duo ring [1, Example 1.1] while $\mathbb{Q}Q_8$ is a duo ring. A natural question which arises is as follows:
 - Question 3.1 Is there any ring R between \mathbb{Z} and \mathbb{Q} (in addition to \mathbb{Q} the field of all rational numbers), such

When is RQ_8 duo?

Group rings over integral domains

- 1 We remark that if G is a non-abelian torsion group and R is a commutative ring with identity for which RG is duo, then RQ_8 is also duo. The question of when RG is duo essentially reduces to that of when RQ_8 is duo, where R is an integral domain.
- 2 As mentioned early, the integral group ring $\mathbb{Z}Q_8$ is a reversible ring, but not a duo ring [1, Example 1.1] while $\mathbb{Q}Q_8$ is a duo ring. A natural question which arises is as follows:
- 3 **Question 3.1** Is there any ring R between \mathbb{Z} and \mathbb{Q} (in addition to \mathbb{Q} the field of all rational numbers), such that RQ_8 is duo.

When is RQ_8 duo?

Group rings over integral domains

- 1 We remark that if G is a non-abelian torsion group and R is a commutative ring with identity for which RG is duo, then RQ_8 is also duo. The question of when RG is duo essentially reduces to that of when RQ_8 is duo, where R is an integral domain.
- 2 As mentioned early, the integral group ring $\mathbb{Z}Q_8$ is a reversible ring, but not a duo ring [1, Example 1.1] while $\mathbb{Q}Q_8$ is a duo ring. A natural question which arises is as follows:
- 3 **Question 3.1** Is there any ring R between \mathbb{Z} and \mathbb{Q} (in addition to \mathbb{Q} the field of all rational numbers), such that RQ_8 is duo.

When is RQ_8 duo?

Group rings over integral domains

- 1 We remark that if G is a non-abelian torsion group and R is a commutative ring with identity for which RG is duo, then RQ_8 is also duo. The question of when RG is duo essentially reduces to that of when RQ_8 is duo, where R is an integral domain.
- 2 As mentioned early, the integral group ring $\mathbb{Z}Q_8$ is a reversible ring, but not a duo ring [1, Example 1.1] while $\mathbb{Q}Q_8$ is a duo ring. A natural question which arises is as follows:
- 3 **Question 3.1** Is there any ring R between \mathbb{Z} and \mathbb{Q} (in addition to \mathbb{Q} the field of all rational numbers), such that RQ_8 is duo.

When is RQ_8 duo?

Group rings over integral domains

- 1 **Theorem 3.5** shows that there does not exist such R . In other words, \mathbb{Q} is the smallest integral domain R of characteristic zero such that RQ_8 is duo.
- We need the following two lemmas. The first is a well known result in number theory and it is a consequence of [6, Theorem 5.14].
- **Lemma 3.2** $1 + x^2 + y^2 \equiv 0 \pmod{p}$ is solvable in \mathbb{Z} for every prime p .
- **Lemma 3.3** Let R be an integral domain such that RQ_8 is duo. If $1 + x^2 + y^2 \neq 0$ for some $x, y \in R$, then $1 + x^2 + y^2$ is invertible in R .



When is RQ_8 duo?

Group rings over integral domains

- 1 Theorem 3.5 shows that there does not exist such R . In other words, \mathbb{Q} is the smallest integral domain R of characteristic zero such that RQ_8 is duo.
- We need the following two lemmas. The first is a well known result in number theory and it is a consequence of [6, Theorem 5.14].
- **Lemma 3.2** $1 + x^2 + y^2 \equiv 0 \pmod{p}$ is solvable in \mathbb{Z} for every prime p .
- **Lemma 3.3** Let R be an integral domain such that RQ_8 is duo. If $1 + x^2 + y^2 \neq 0$ for some $x, y \in R$, then $1 + x^2 + y^2$ is invertible in R .



When is RQ_8 duo?

Group rings over integral domains

- 1 Theorem 3.5 shows that there does not exist such R . In other words, \mathbb{Q} is the smallest integral domain R of characteristic zero such that RQ_8 is duo.
- 2 We need the following two lemmas. The first is a well known result in number theory and it is a consequence of [6, Theorem 5.14].
- 3 **Lemma 3.2** $1 + x^2 + y^2 \equiv 0 \pmod{p}$ is solvable in \mathbb{Z} for every prime p .
- 4 **Lemma 3.3** Let R be an integral domain such that RQ_8 is duo. If $1 + x^2 + y^2 \neq 0$ for some $x, y \in R$, then $1 + x^2 + y^2$ is invertible in R .

When is RQ_8 duo?

Group rings over integral domains

- 1 Theorem 3.5 shows that there does not exist such R . In other words, \mathbb{Q} is the smallest integral domain R of characteristic zero such that RQ_8 is duo.
- 2 We need the following two lemmas. The first is a well known result in number theory and it is a consequence of [6, Theorem 5.14].
 - 3 **Lemma 3.2** $1 + x^2 + y^2 \equiv 0 \pmod{p}$ is solvable in \mathbb{Z} for every prime p .
 - 1 **Lemma 3.3** Let R be an integral domain such that RQ_8 is duo. If $1 + x^2 + y^2 \neq 0$ for some $x, y \in R$, then $1 + x^2 + y^2$ is invertible in R .

When is RQ_8 duo?

Group rings over integral domains

- 1 Theorem 3.5 shows that there does not exist such R . In other words, \mathbb{Q} is the smallest integral domain R of characteristic zero such that RQ_8 is duo.
- 2 We need the following two lemmas. The first is a well known result in number theory and it is a consequence of [6, Theorem 5.14].
- 3 **Lemma 3.2** $1 + x^2 + y^2 \equiv 0 \pmod{p}$ is solvable in \mathbb{Z} for every prime p .
- 4 **Lemma 3.3** Let R be an integral domain such that RQ_8 is duo. If $1 + x^2 + y^2 \neq 0$ for some $x, y \in R$, then $1 + x^2 + y^2$ is invertible in R .

When is RQ_8 duo?

Group rings over integral domains

- 1 Theorem 3.5 shows that there does not exist such R . In other words, \mathbb{Q} is the smallest integral domain R of characteristic zero such that RQ_8 is duo.
- 2 We need the following two lemmas. The first is a well known result in number theory and it is a consequence of [6, Theorem 5.14].
- 3 **Lemma 3.2** $1 + x^2 + y^2 \equiv 0 \pmod{p}$ is solvable in \mathbb{Z} for every prime p .
- 4 **Lemma 3.3** Let R be an integral domain such that RQ_8 is duo. If $1 + x^2 + y^2 \neq 0$ for some $x, y \in R$, then $1 + x^2 + y^2$ is invertible in R .

When is RQ_8 duo?

Group rings over integral domains

- 1 Theorem 3.5 shows that there does not exist such R . In other words, \mathbb{Q} is the smallest integral domain R of characteristic zero such that RQ_8 is duo.
- 2 We need the following two lemmas. The first is a well known result in number theory and it is a consequence of [6, Theorem 5.14].
- 3 **Lemma 3.2** $1 + x^2 + y^2 \equiv 0 \pmod{p}$ is solvable in \mathbb{Z} for every prime p .
- 4 **Lemma 3.3** Let R be an integral domain such that RQ_8 is duo. If $1 + x^2 + y^2 \neq 0$ for some $x, y \in R$, then $1 + x^2 + y^2$ is invertible in R .

When is RQ_8 duo?

Group rings over integral domains

- ① **Proof:** If R is finite, then R is a field, and thus the result holds. Next we assume that R is infinite.
- ② We first show that if $1 + x_0^2 \neq 0$ for some $x_0 \in R$, then $1 + x_0^2$ is invertible in R .
- ③ We then show that if $(1 + x^2 + y^2) \neq 0$ for some $x, y \in R$, then it is a factor of $1 + x_0^2 (\neq 0)$ for some $x_0 \in R$, so, it is invertible in R .
- ④ Note that if R is an integral domain such that RQ_8 is duo, then RQ_8 is reversible. It follows from [4, Theorem 2.5] that the characteristic of R is either 2 or 0. In the latter case, by [4, Theorem 4.2] (see also [3, Theorem 3.1]), we have $1 + x^2 + y^2 \neq 0$, for all $x, y \in R$. As a consequence of the above theorem, we obtain



When is RQ_8 duo?

Group rings over integral domains

- 1 **Proof:** If R is finite, then R is a field, and thus the result holds. Next we assume that R is infinite.
- 2 **We first show that if $1 + x_0^2 \neq 0$ for some $x_0 \in R$, then $1 + x_0^2$ is invertible in R .**
- 3 We then show that if $(1 + x^2 + y^2) \neq 0$ for some $x, y \in R$, then it is a factor of $1 + x_0^2 (\neq 0)$ for some $x_0 \in R$, so, it is invertible in R .
- 4 Note that if R is an integral domain such that RQ_8 is duo, then RQ_8 is reversible. It follows from [4, Theorem 2.5] that the characteristic of R is either 2 or 0. In the latter case, by [4, Theorem 4.2] (see also [3, Theorem 3.1]), we have $1 + x^2 + y^2 \neq 0$, for all $x, y \in R$. As a consequence, RQ_8 is reversible and duo.



When is RQ_8 duo?

Group rings over integral domains

- 1 **Proof:** If R is finite, then R is a field, and thus the result holds. Next we assume that R is infinite.
- 2 We first show that if $1 + x_0^2 \neq 0$ for some $x_0 \in R$, then $1 + x_0^2$ is invertible in R .
- 3 We then show that if $(1 + x^2 + y^2) \neq 0$ for some $x, y \in R$, then it is a factor of $1 + x_0^2 (\neq 0)$ for some $x_0 \in R$, so, it is invertible in R .
- 4 Note that if R is an integral domain such that RQ_8 is duo, then RQ_8 is reversible. It follows from [4, Theorem 2.5] that the characteristic of R is either 2 or 0. In the latter case, by [4, Theorem 4.2] (see also [3, Theorem 3.1]), we have $1 + x^2 + y^2 \neq 0$, for all $x, y \in R$. As a

When is RQ_8 duo?

Group rings over integral domains

- 1 **Proof:** If R is finite, then R is a field, and thus the result holds. Next we assume that R is infinite.
- 2 We first show that if $1 + x_0^2 \neq 0$ for some $x_0 \in R$, then $1 + x_0^2$ is invertible in R .
- 3 We then show that if $(1 + x^2 + y^2) \neq 0$ for some $x, y \in R$, then it is a factor of $1 + x_0^2 (\neq 0)$ for some $x_0 \in R$, so, it is invertible in R .
- 4 **Note that if R is an integral domain such that RQ_8 is duo, then RQ_8 is reversible.** It follows from [4, Theorem 2.5] that the characteristic of R is either 2 or 0. In the latter case, by [4, Theorem 4.2] (see also [3, Theorem 3.1]), we have $1 + x^2 + y^2 \neq 0$, for all $x, y \in R$. As a consequence of the above lemma, we obtain

When is RQ_8 duo?

Group rings over integral domains

- 1 **Proof:** If R is finite, then R is a field, and thus the result holds. Next we assume that R is infinite.
- 2 We first show that if $1 + x_0^2 \neq 0$ for some $x_0 \in R$, then $1 + x_0^2$ is invertible in R .
- 3 We then show that if $(1 + x^2 + y^2) \neq 0$ for some $x, y \in R$, then it is a factor of $1 + x_0^2 (\neq 0)$ for some $x_0 \in R$, so, it is invertible in R .
- 4 Note that if R is an integral domain such that RQ_8 is duo, then RQ_8 is reversible. It follows from [4, Theorem 2.5] that the characteristic of R is either 2 or 0. In the latter case, by [4, Theorem 4.2] (see also [3, Theorem 3.1]), we have $1 + x^2 + y^2 \neq 0$, for all $x, y \in R$. As a consequence of the above lemma, we obtain

When is RQ_8 duo?

Group rings over integral domains

- 1 **Proof:** If R is finite, then R is a field, and thus the result holds. Next we assume that R is infinite.
- 2 We first show that if $1 + x_0^2 \neq 0$ for some $x_0 \in R$, then $1 + x_0^2$ is invertible in R .
- 3 We then show that if $(1 + x^2 + y^2) \neq 0$ for some $x, y \in R$, then it is a factor of $1 + x_0^2 (\neq 0)$ for some $x_0 \in R$, so, it is invertible in R .
- 4 Note that if R is an integral domain such that RQ_8 is duo, then RQ_8 is reversible. It follows from [4, Theorem 2.5] that the characteristic of R is either 2 or 0. In the latter case, by [4, Theorem 4.2] (see also [3, Theorem 3.1]), we have $1 + x^2 + y^2 \neq 0$, for all $x, y \in R$. As a consequence of the above lemma, we obtain

When is RQ_8 duo?

Group rings over integral domains

- 1 **Proof:** If R is finite, then R is a field, and thus the result holds. Next we assume that R is infinite.
- 2 We first show that if $1 + x_0^2 \neq 0$ for some $x_0 \in R$, then $1 + x_0^2$ is invertible in R .
- 3 We then show that if $(1 + x^2 + y^2) \neq 0$ for some $x, y \in R$, then it is a factor of $1 + x_0^2 (\neq 0)$ for some $x_0 \in R$, so, it is invertible in R .
- 4 Note that if R is an integral domain such that RQ_8 is duo, then RQ_8 is reversible. It follows from [4, Theorem 2.5] that the characteristic of R is either 2 or 0. In the latter case, by [4, Theorem 4.2] (see also [3, Theorem 3.1]), we have $1 + x^2 + y^2 \neq 0$, for all $x, y \in R$. As a consequence of the above lemma, we obtain

When is RQ_8 duo?

Group rings over integral domains

- 1 **Proof:** If R is finite, then R is a field, and thus the result holds. Next we assume that R is infinite.
- 2 We first show that if $1 + x_0^2 \neq 0$ for some $x_0 \in R$, then $1 + x_0^2$ is invertible in R .
- 3 We then show that if $(1 + x^2 + y^2) \neq 0$ for some $x, y \in R$, then it is a factor of $1 + x_0^2 (\neq 0)$ for some $x_0 \in R$, so, it is invertible in R .
- 4 Note that if R is an integral domain such that RQ_8 is duo, then RQ_8 is reversible. It follows from [4, Theorem 2.5] that the characteristic of R is either 2 or 0. In the latter case, by [4, Theorem 4.2] (see also [3, Theorem 3.1]), we have $1 + x^2 + y^2 \neq 0$, for all $x, y \in R$. As a consequence of the above lemma, we obtain

When is RQ_8 duo?

Group rings over integral domains

- 1 **Proof:** If R is finite, then R is a field, and thus the result holds. Next we assume that R is infinite.
- 2 We first show that if $1 + x_0^2 \neq 0$ for some $x_0 \in R$, then $1 + x_0^2$ is invertible in R .
- 3 We then show that if $(1 + x^2 + y^2) \neq 0$ for some $x, y \in R$, then it is a factor of $1 + x_0^2 (\neq 0)$ for some $x_0 \in R$, so, it is invertible in R .
- 4 Note that if R is an integral domain such that RQ_8 is duo, then RQ_8 is reversible. It follows from [4, Theorem 2.5] that the characteristic of R is either 2 or 0. In the latter case, by [4, Theorem 4.2] (see also [3, Theorem 3.1]), we have $1 + x^2 + y^2 \neq 0$, for all $x, y \in R$. As a consequence of the above lemma, we obtain

When is RQ_8 duo?

Group rings over integral domains

- **Corollary 3.4** Let R be an integral domain such that RQ_8 is duo. Then $\text{char } R = 2$ or $\text{char } R = 0$. In the latter case, we have $1 + x^2 + y^2 \in \mathcal{U}(R)$, for all $x, y \in R$.
- Theorem 3.5 [2, Theorem 2.4] Let R be an integral domain such that RQ_8 is duo. Then the following statements hold.
 - (1) If $\text{char}(R) \neq 0$, then R must be a field.
 - (2) If S is a ring of algebraic integers with its quotient field K_S such that $S \subseteq R \subseteq K_S$, then $R = K_S$. In particular, if $\mathbb{Z} \subseteq R \subseteq \mathbb{Q}$, then $R = \mathbb{Q}$.
- **Proof:** (1) We note that $\text{char}(R) = 2$ by Corollary 4. Let $\alpha \neq 0 \in R$ and $x = \alpha - 1 \in R$. Then $1 + x^2 = (1 + x)^2 = \alpha^2 \neq 0$. It follows from Lemma 4 that α^2 is invertible in R and so is α . Therefore, R is a field.



When is RQ_8 duo?

Group rings over integral domains

- **Corollary 3.4** Let R be an integral domain such that RQ_8 is duo. Then $\text{char } R = 2$ or $\text{char } R = 0$. In the latter case, we have $1 + x^2 + y^2 \in \mathcal{U}(R)$, for all $x, y \in R$.
- **Theorem 3.5** [2, Theorem 2.4] Let R be an integral domain such that RQ_8 is duo. Then the following statements hold.
 - (1) If $\text{char}(R) \neq 0$, then R must be a field.
 - (2) If S is a ring of algebraic integers with its quotient field K_S such that $S \subseteq R \subseteq K_S$, then $R = K_S$. In particular, if $\mathbb{Z} \subseteq R \subseteq \mathbb{Q}$, then $R = \mathbb{Q}$.
- **Proof:** (1) We note that $\text{char}(R) = 2$ by Corollary 4. Let $\alpha \neq 0 \in R$ and $x = \alpha - 1 \in R$. Then $1 + x^2 = (1 + x)^2 = \alpha^2 \neq 0$. It follows from Lemma 4 that α^2 is invertible in R and so is α . Therefore, R is a field.

When is RQ_8 duo?

Group rings over integral domains

- **Corollary 3.4** Let R be an integral domain such that RQ_8 is duo. Then $\text{char } R = 2$ or $\text{char } R = 0$. In the latter case, we have $1 + x^2 + y^2 \in \mathcal{U}(R)$, for all $x, y \in R$.
- **Theorem 3.5** [2, Theorem 2.4] Let R be an integral domain such that RQ_8 is duo. Then the following statements hold.

(1) If $\text{char}(R) \neq 0$, then R must be a field.

If S is a ring of algebraic integers with its quotient field K_S such that $S \subseteq R \subseteq K_S$, then $R = K_S$. In particular, if $\mathbb{Z} \subseteq R \subseteq \mathbb{Q}$, then $R = \mathbb{Q}$.

- **Proof:** (1) We note that $\text{char}(R) = 2$ by Corollary 4. Let $\alpha \neq 0 \in R$ and $x = \alpha - 1 \in R$. Then $1 + x^2 = (1 + x)^2 = \alpha^2 \neq 0$. It follows from Lemma 4 that α^2 is invertible in R and so is α . Therefore, R is a field.

When is RQ_8 duo?

Group rings over integral domains

- **Corollary 3.4** Let R be an integral domain such that RQ_8 is duo. Then $\text{char } R = 2$ or $\text{char } R = 0$. In the latter case, we have $1 + x^2 + y^2 \in \mathcal{U}(R)$, for all $x, y \in R$.
- **Theorem 3.5** [2, Theorem 2.4] Let R be an integral domain such that RQ_8 is duo. Then the following statements hold.
 - (1) If $\text{char}(R) \neq 0$, then R must be a field.
 - (2) If S is a ring of algebraic integers with its quotient field K_S such that $S \subseteq R \subseteq K_S$, then $R = K_S$. In particular, if $\mathbb{Z} \subseteq R \subseteq \mathbb{Q}$, then $R = \mathbb{Q}$.
- **Proof:** (1) We note that $\text{char}(R) = 2$ by Corollary 4. Let $\alpha \neq 0 \in R$ and $x = \alpha - 1 \in R$. Then $1 + x^2 = (1 + x)^2 = \alpha^2 \neq 0$. It follows from Lemma 4 that α^2 is invertible in R and so is α . Therefore, R is a field.

When is RQ_8 duo?

Group rings over integral domains

- Corollary 3.4** Let R be an integral domain such that RQ_8 is duo. Then $\text{char } R = 2$ or $\text{char } R = 0$. In the latter case, we have $1 + x^2 + y^2 \in \mathcal{U}(R)$, for all $x, y \in R$.
- Theorem 3.5** [2, Theorem 2.4] Let R be an integral domain such that RQ_8 is duo. Then the following statements hold.
 - (1) If $\text{char}(R) \neq 0$, then R must be a field.
 - (2) If S is a ring of algebraic integers with its quotient field K_S such that $S \subseteq R \subseteq K_S$, then $R = K_S$. In particular, if $\mathbb{Z} \subseteq R \subseteq \mathbb{Q}$, then $R = \mathbb{Q}$.
- Proof:** (1) We note that $\text{char}(R) = 2$ by Corollary 4. Let $\alpha \neq 0 \in R$ and $x = \alpha - 1 \in R$. Then $1 + x^2 = (1 + x)^2 = \alpha^2 \neq 0$. It follows from Lemma 4 that α^2 is invertible in R and so is α . Therefore, R is a field.

When is RQ_8 duo?

Group rings over integral domains

- **Corollary 3.4** Let R be an integral domain such that RQ_8 is duo. Then $\text{char } R = 2$ or $\text{char } R = 0$. In the latter case, we have $1 + x^2 + y^2 \in \mathcal{U}(R)$, for all $x, y \in R$.
- **Theorem 3.5** [2, Theorem 2.4] Let R be an integral domain such that RQ_8 is duo. Then the following statements hold.
 - (1) If $\text{char}(R) \neq 0$, then R must be a field.
 - (2) If S is a ring of algebraic integers with its quotient field K_S such that $S \subseteq R \subseteq K_S$, then $R = K_S$. In particular, if $\mathbb{Z} \subseteq R \subseteq \mathbb{Q}$, then $R = \mathbb{Q}$.
- **Proof:** (1) We note that $\text{char}(R) = 2$ by Corollary 4. Let $\alpha \neq 0 \in R$ and $x = \alpha - 1 \in R$. Then $1 + x^2 = (1 + x)^2 = \alpha^2 \neq 0$. It follows from Lemma 4 that α^2 is invertible in R and so is α . Therefore, R is a field.

When is RQ_8 duo?

Group rings over integral domains

- **Corollary 3.4** Let R be an integral domain such that RQ_8 is duo. Then $\text{char } R = 2$ or $\text{char } R = 0$. In the latter case, we have $1 + x^2 + y^2 \in \mathcal{U}(R)$, for all $x, y \in R$.
- **Theorem 3.5** [2, Theorem 2.4] Let R be an integral domain such that RQ_8 is duo. Then the following statements hold.
 - (1) If $\text{char}(R) \neq 0$, then R must be a field.
 - (2) If S is a ring of algebraic integers with its quotient field K_S such that $S \subseteq R \subseteq K_S$, then $R = K_S$. In particular, if $\mathbb{Z} \subseteq R \subseteq \mathbb{Q}$, then $R = \mathbb{Q}$.
- **Proof:** (1) We note that $\text{char}(R) = 2$ by Corollary 4. Let $\alpha \neq 0 \in R$ and $x = \alpha - 1 \in R$. Then $1 + x^2 = (1 + x)^2 = \alpha^2 \neq 0$. It follows from Lemma 4 that α^2 is invertible in R and so is α . Therefore, R is a field.

When is RQ_8 duo?

Group rings over integral domains

- (2) It suffices to prove that every nonzero element $\alpha \in S$ is invertible in R , and thus $K_S \subseteq R$. We first prove that if $0 \neq \alpha \in \mathbb{Z}$, then α is invertible in R . Let p be any prime. By Lemma 3.2, $p \mid 1 + x^2 + y^2$ for some $x, y \in \mathbb{Z}$. It follows from Corollary 3.4 that $1 + x^2 + y^2$, and thus p is invertible in R . Therefore, α is invertible in R .
- Let $0 \neq \alpha \in S$. Then there is a monic polynomial $f(x) \in \mathbb{Z}[x]$ such that $f(\alpha) = 0$, where

$$f(x) = x^n + c_{n-1}x^{n-1} + \cdots + c_1x + c_0,$$

with all $c_i \in \mathbb{Z}$ and $c_0 \neq 0$. Then

$$(\alpha^{n-1} + c_{n-1}\alpha^{n-2} + \cdots + c_1)\alpha = -c_0.$$

Since $-c_0$ is invertible in R , α is also invertible in R .

When is RQ_8 duo?

Group rings over integral domains

- (2) It suffices to prove that every nonzero element $\alpha \in S$ is invertible in R , and thus $K_S \subseteq R$. We first prove that if $0 \neq \alpha \in \mathbb{Z}$, then α is invertible in R . Let p be any prime. By Lemma 3.2, $p \mid 1 + x^2 + y^2$ for some $x, y \in \mathbb{Z}$. It follows from Corollary 3.4 that $1 + x^2 + y^2$, and thus p is invertible in R . Therefore, α is invertible in R .
- **Let $0 \neq \alpha \in S$. Then there is a monic polynomial $f(x) \in \mathbb{Z}[x]$ such that $f(\alpha) = 0$, where**

$$f(x) = x^n + c_{n-1}x^{n-1} + \cdots + c_1x + c_0,$$

with all $c_i \in \mathbb{Z}$ and $c_0 \neq 0$. Then

$$(\alpha^{n-1} + c_{n-1}\alpha^{n-2} + \cdots + c_1)\alpha = -c_0.$$

Since $-c_0$ is invertible in R , α is also invertible in R .

When is RQ_8 duo?

Group rings over integral domains

- (2) It suffices to prove that every nonzero element $\alpha \in S$ is invertible in R , and thus $K_S \subseteq R$. We first prove that if $0 \neq \alpha \in \mathbb{Z}$, then α is invertible in R . Let p be any prime. By Lemma 3.2, $p | 1 + x^2 + y^2$ for some $x, y \in \mathbb{Z}$. It follows from Corollary 3.4 that $1 + x^2 + y^2$, and thus p is invertible in R . Therefore, α is invertible in R .
- Let $0 \neq \alpha \in S$. Then there is a monic polynomial $f(x) \in \mathbb{Z}[x]$ such that $f(\alpha) = 0$, where

$$f(x) = x^n + c_{n-1}x^{n-1} + \cdots + c_1x + c_0,$$

with all $c_i \in \mathbb{Z}$ and $c_0 \neq 0$. Then

$$(\alpha^{n-1} + c_{n-1}\alpha^{n-2} + \cdots + c_1)\alpha = -c_0.$$

Since $-c_0$ is invertible in R , α is also invertible in R .

When is RQ_8 duo?

Group rings over integral domains

- (2) It suffices to prove that every nonzero element $\alpha \in S$ is invertible in R , and thus $K_S \subseteq R$. We first prove that if $0 \neq \alpha \in \mathbb{Z}$, then α is invertible in R . Let p be any prime. By Lemma 3.2, $p|1 + x^2 + y^2$ for some $x, y \in \mathbb{Z}$. It follows from Corollary 3.4 that $1 + x^2 + y^2$, and thus p is invertible in R . Therefore, α is invertible in R .
- Let $0 \neq \alpha \in S$. Then there is a monic polynomial $f(x) \in \mathbb{Z}[x]$ such that $f(\alpha) = 0$, where

$$f(x) = x^n + c_{n-1}x^{n-1} + \cdots + c_1x + c_0,$$

with all $c_i \in \mathbb{Z}$ and $c_0 \neq 0$. Then

$$(\alpha^{n-1} + c_{n-1}\alpha^{n-2} + \cdots + c_1)\alpha = -c_0.$$

Since $-c_0$ is invertible in R , α is also invertible in R .

When is RQ_8 duo?

Group rings over integral domains

- 1 It appears that Theorem 3.5 suggests that if RQ_8 is duo, then R is a field. However, the following proposition shows that this is not the case.
- 2 Proposition 3.6 Let $S = \mathbb{Q}[x]$ be the polynomial ring over the rational field, and S_P be the localization of S at the maximal ideal $P = \langle x \rangle$. Then $R = S_P$ is a local integral domain of characteristic 0, but not a field, such that RQ_8 is duo.



When is RQ_8 duo?

Group rings over integral domains

- 1 It appears that Theorem 3.5 suggests that if RQ_8 is duo, then R is a field. However, the following proposition shows that this is not the case.
- 2 **Proposition 3.6** Let $S = \mathbb{Q}[x]$ be the polynomial ring over the rational field, and S_P be the localization of S at the maximal ideal $P = \langle x \rangle$. Then $R = S_P$ is a local integral domain of characteristic 0, but not a field, such that RQ_8 is duo.



When is RQ_8 duo?

Group rings over integral domains

- 1 It appears that Theorem 3.5 suggests that if RQ_8 is duo, then R is a field. However, the following proposition shows that this is not the case.
- 2 Proposition 3.6 Let $S = \mathbb{Q}[x]$ be the polynomial ring over the rational field, and S_P be the localization of S at the maximal ideal $P = \langle x \rangle$. Then $R = S_P$ is a local integral domain of characteristic 0, but not a field, such that RQ_8 is duo.

When is RQ_8 duo?

Group rings over integral domains

- 1 It appears that Theorem 3.5 suggests that if RQ_8 is duo, then R is a field. However, the following proposition shows that this is not the case.
- 2 Proposition 3.6 Let $S = \mathbb{Q}[x]$ be the polynomial ring over the rational field, and S_P be the localization of S at the maximal ideal $P = \langle x \rangle$. Then $R = S_P$ is a local integral domain of characteristic 0, but not a field, such that RQ_8 is duo.

When is RG duo?

Group rings over integral domains

- ① Theorem 3.5 together with Theorem 2.5 gives a description of when RG is duo.
- ② Theorem 3.7 If R is an integral domain of $\text{char}(R) \neq 0$ and G is a non-abelian torsion group, then the following statements are equivalent:
 - (1) RG is duo.
 - (2) R is a field and RG is reversible.
 - (3) $G = Q_8 \times E'_2$, $R = K$ is a field of characteristic 2 and the equation $1 + x + x^2 = 0$ has no solutions in any cyclotomic field $K(\xi_d)$ for any odd d which is an order of an element of E'_2 .



When is RG duo?

Group rings over integral domains

- 1 Theorem 3.5 together with Theorem 2.5 gives a description of when RG is duo.
- 2 Theorem 3.7 If R is an integral domain of $\text{char}(R) \neq 0$ and G is a non-abelian torsion group, then the following statements are equivalent:
 - (1) RG is duo.
 - (2) R is a field and RG is reversible.
 - (3) $G = Q_8 \times E_2^d$, $R = K$ is a field of characteristic 2 and the equation $1 + x + x^2 = 0$ has no solutions in any cyclotomic field $K(\xi_d)$ for any odd d which is an order of an element of E_2^d .



When is RG duo?

Group rings over integral domains

- 1 Theorem 3.5 together with Theorem 2.5 gives a description of when RG is duo.
- 2 Theorem 3.7 If R is an integral domain of $\text{char}(R) \neq 0$ and G is a non-abelian torsion group, then the following statements are equivalent:
 - (1) RG is duo.
 - (2) R is a field and RG is reversible.
 - (3) $G = Q_8 \times E'_2$, $R = K$ is a field of characteristic 2 and the equation $1 + x + x^2 = 0$ has no solutions in any cyclotomic field $K(\xi_d)$ for any odd d which is an order of an element of E'_2 .



When is RG duo?

Group rings over integral domains

- 1 Theorem 3.5 together with Theorem 2.5 gives a description of when RG is duo.
- 2 Theorem 3.7 If R is an integral domain of $\text{char}(R) \neq 0$ and G is a non-abelian torsion group, then the following statements are equivalent:
 - (1) RG is duo.
 - (2) R is a field and RG is reversible.
 - (3) $G = Q_8 \times E'_2$, $R = K$ is a field of characteristic 2 and the equation $1 + x + x^2 = 0$ has no solutions in any cyclotomic field $K(\xi_d)$ for any odd d which is an order of an element of E'_2 .



When is RG duo?

Group rings over integral domains

- 1 Theorem 3.5 together with Theorem 2.5 gives a description of when RG is duo.
- 2 Theorem 3.7 If R is an integral domain of $\text{char}(R) \neq 0$ and G is a non-abelian torsion group, then the following statements are equivalent:
 - (1) RG is duo.
 - (2) R is a field and RG is reversible.
 - (3) $G = Q_8 \times E'_2$, $R = K$ is a field of characteristic 2 and the equation $1 + x + x^2 = 0$ has no solutions in any cyclotomic field $K(\xi_d)$ for any odd d which is an order of an element of E'_2 .

When is RG duo?

Group rings over integral domains

- ① Theorem 3.5 together with Theorem 2.5 gives a description of when RG is duo.
- ② Theorem 3.7 If R is an integral domain of $\text{char}(R) \neq 0$ and G is a non-abelian torsion group, then the following statements are equivalent:
 - (1) RG is duo.
 - (2) R is a field and RG is reversible.
 - (3) $G = Q_8 \times E'_2$, $R = K$ is a field of characteristic 2 and the equation $1 + x + x^2 = 0$ has no solutions in any cyclotomic field $K(\xi_d)$ for any odd d which is an order of an element of E'_2 .

When is RG duo?

Group rings over integral domains

- 1 Theorem 3.5 together with Theorem 2.5 gives a description of when RG is duo.
- 2 Theorem 3.7 If R is an integral domain of $\text{char}(R) \neq 0$ and G is a non-abelian torsion group, then the following statements are equivalent:
 - (1) RG is duo.
 - (2) R is a field and RG is reversible.
 - (3) $G = Q_8 \times E'_2$, $R = K$ is a field of characteristic 2 and the equation $1 + x + x^2 = 0$ has no solutions in any cyclotomic field $K(\xi_d)$ for any odd d which is an order of an element of E'_2 .

When is RG duo?

Group rings over integral domains

- ① **Theorem 3.8** If R is an integral domain of $\text{char}(R) = 0$ such that $S \subseteq R \subseteq K_S$, and G is a non-abelian torsion group, then the following statements are equivalent:
- (1) RG is duo.
 - (2) R is a field and RG is reversible.
 - (3) $G = Q_8 \times E_2 \times E'_3$, $R = K$ is a field of characteristic 0 and the equation $1 + x^2 + y^2 = 0$ has no solutions in any cyclotomic field $K(\zeta_d)$ for any odd d which is an order of an element of E'_3 .



When is RG duo?

Group rings over integral domains

- ① Theorem 3.8 If R is an integral domain of $\text{char}(R) = 0$ such that $S \subseteq R \subseteq K_S$, and G is a non-abelian torsion group, then the following statements are equivalent:

(1) RG is duo.

R is a field and RG is reversible.

- (2) $G = Q_8 \times E_2 \times E'_2$, $R = K$ is a field of characteristic 0 and the equation $1 + x^2 + y^2 = 0$ has no solutions in any cyclotomic field $K(\xi_d)$ for any odd d which is an order of an element of E'_2 .



When is RG duo?

Group rings over integral domains

- ① Theorem 3.8 If R is an integral domain of $\text{char}(R) = 0$ such that $S \subseteq R \subseteq K_S$, and G is a non-abelian torsion group, then the following statements are equivalent:

(1) RG is duo.

(2) R is a field and RG is reversible.

(3) $G = Q_8 \times E_2 \times E'_2$, $R = K$ is a field of characteristic 0 and the equation $1 + x^2 + y^2 = 0$ has no solutions in any cyclotomic field $K(\xi_d)$ for any odd d which is an order of an element of E'_2 .



When is RG duo?

Group rings over integral domains

- ① Theorem 3.8 If R is an integral domain of $\text{char}(R) = 0$ such that $S \subseteq R \subseteq K_S$, and G is a non-abelian torsion group, then the following statements are equivalent:
- (1) RG is duo.
 - (2) R is a field and RG is reversible.
 - (3) $G = Q_8 \times E_2 \times E'_2$, $R = K$ is a field of characteristic 0 and the equation $1 + x^2 + y^2 = 0$ has no solutions in any cyclotomic field $K(\xi_d)$ for any odd d which is an order of an element of E'_2 .



When is RG duo?

Group rings over integral domains

- ① Theorem 3.8 If R is an integral domain of $\text{char}(R) = 0$ such that $S \subseteq R \subseteq K_S$, and G is a non-abelian torsion group, then the following statements are equivalent:
- (1) RG is duo.
 - (2) R is a field and RG is reversible.
 - (3) $G = Q_8 \times E_2 \times E'_2$, $R = K$ is a field of characteristic 0 and the equation $1 + x^2 + y^2 = 0$ has no solutions in any cyclotomic field $K(\xi_d)$ for any odd d which is an order of an element of E'_2 .



When is RG duo?

Group rings over integral domains

- ① Theorem 3.8 If R is an integral domain of $\text{char}(R) = 0$ such that $S \subseteq R \subseteq K_S$, and G is a non-abelian torsion group, then the following statements are equivalent:
- (1) RG is duo.
 - (2) R is a field and RG is reversible.
 - (3) $G = Q_8 \times E_2 \times E'_2$, $R = K$ is a field of characteristic 0 and the equation $1 + x^2 + y^2 = 0$ has no solutions in any cyclotomic field $K(\xi_d)$ for any odd d which is an order of an element of E'_2 .



When is RQ duo?

Group rings over integral domains

- 1 Recall that Proposition 3.6 indicates that there exists an integral domain R , which is not a field, such that RQ_8 is duo. In that case, a necessary condition for RQ_8 to be duo is given in Corollary 3.4, i.e. $1+x^2+y^2 \in \mathcal{U}(R)$ for all $x, y \in R$. We are not aware of any example of an integral domain R of $\text{char}(R) = 0$ satisfying this necessary condition for which RQ_8 is not duo. We close this talk by proposing the following question.
- Question 3.10 Assume that R is an integral domain of $\text{char}(R) = 0$ such that $1+x^2+y^2 \in \mathcal{U}(R)$ for all $x, y \in R$. Is RQ_8 duo?



When is RQ duo?

Group rings over integral domains

- 1 Recall that Proposition 3.6 indicates that there exists an integral domain R , which is not a field, such that RQ_8 is duo. In that case, a necessary condition for RQ_8 to be duo is given in Corollary 3.4, i.e. $1+x^2+y^2 \in \mathcal{U}(R)$ for all $x, y \in R$. We are not aware of any example of an integral domain R of $\text{char}(R) = 0$ satisfying this necessary condition for which RQ_8 is not duo. We close this talk by proposing the following question.
- Question 3.10 Assume that R is an integral domain of $\text{char}(R) = 0$ such that $1+x^2+y^2 \in \mathcal{U}(R)$ for all $x, y \in R$. Is RQ_8 duo?



When is RQ_8 duo?

Group rings over integral domains

- 1 Recall that Proposition 3.6 indicates that there exists an integral domain R , which is not a field, such that RQ_8 is duo. In that case, a necessary condition for RQ_8 to be duo is given in Corollary 3.4, i.e. $1+x^2+y^2 \in \mathcal{U}(R)$ for all $x, y \in R$. We are not aware of any example of an integral domain R of $\text{char}(R) = 0$ satisfying this necessary condition for which RQ_8 is not duo. We close this talk by proposing the following question.
- 2 **Question 3.10** Assume that R is an integral domain of $\text{char}(R) = 0$ such that $1+x^2+y^2 \in \mathcal{U}(R)$ for all $x, y \in R$. Is RQ_8 duo?

When is RQ duo?

Group rings over integral domains

- 1 Recall that Proposition 3.6 indicates that there exists an integral domain R , which is not a field, such that RQ_8 is duo. In that case, a necessary condition for RQ_8 to be duo is given in Corollary 3.4, i.e. $1+x^2+y^2 \in \mathcal{U}(R)$ for all $x, y \in R$. We are not aware of any example of an integral domain R of $\text{char}(R) = 0$ satisfying this necessary condition for which RQ_8 is not duo. We close this talk by proposing the following question.
- 2 **Question 3.10** Assume that R is an integral domain of $\text{char}(R) = 0$ such that $1+x^2+y^2 \in \mathcal{U}(R)$ for all $x, y \in R$. Is RQ_8 duo?

When is RQ duo?

Group rings over integral domains

- 1 Recall that Proposition 3.6 indicates that there exists an integral domain R , which is not a field, such that RQ_8 is duo. In that case, a necessary condition for RQ_8 to be duo is given in Corollary 3.4, i.e. $1+x^2+y^2 \in \mathcal{U}(R)$ for all $x, y \in R$. We are not aware of any example of an integral domain R of $\text{char}(R) = 0$ satisfying this necessary condition for which RQ_8 is not duo. We close this talk by proposing the following question.
- 2 **Question 3.10** Assume that R is an integral domain of $\text{char}(R) = 0$ such that $1+x^2+y^2 \in \mathcal{U}(R)$ for all $x, y \in R$. Is RQ_8 duo?

References

-  H. Bell and Y. Li, Duo group rings, *J. Pure Appl. Algebra*, **209** (2007), 833 - 838.
-  W. Gao and Y. Li, On Duo group rings, *Algebra Colloq.*, To Appear.
-  M. Gutan and A. Kisielewicz, *Reversible group rings*, *J. Algebra* **279** (2004), 280–291.
-  Y. Li and M.M. Parmenter, Reversible group rings over commutative rings, *Comm. Algebra* **35** (2007), 4096–4104.
-  G. Marks, *Reversible and symmetric rings*, *J. Pure Appl. Algebra* **174** (2002), 311 - 318.
-  I. Niven, H.S. Zuckerman and H.L. Montgomery, *An introduction to the theory of numbers*, Fifth edition, John Wiley & Sons, Inc., New York, 1991.
-  C. Polcino Milies and S.K. Sehgal, *An Introduction to Group Rings*, Kluwer Academic Publishers, Dordrecht, 2002.

Thank you!

AUTHOR: Yuanlin Li

ADDRESS: Department of Mathematics
Brock University
Canada

EMAIL: yli@brocku.ca

