

$${}^2A_6(q) \bmod l, d = 10$$

	<i>ps</i>	<i>ps</i>	11111	21	2111	<i>ps</i>	<i>ps</i>	11111	21	2111	<i>ps</i>	<i>ps</i>	<i>ps</i>	<i>ps</i>	<i>ps</i>
3.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.
1.2	1	1	.	.	.	.	.	.	.	.	.	.	.	.	.
.21	.	1	1	.	.	.	.	.	.	.	.	.	.	.	.
${}^2A_2 : 11.$	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.
${}^2A_2 : .11$	.	.	.	1	1	.	.	.	.	.	.	.	.	.	.
21.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.
11.1	.	.	.	.	.	1	1	.	.	.	.	.	.	.	.
.111	.	.	.	.	.	.	1	1	.	.	.	.	.	.	.
${}^2A_2 : 2.$	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.
${}^2A_2 : .2$	.	.	.	.	.	.	.	.	1	1	.	.	.	.	.
2.1	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.
.3	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.
${}^2A_2 : 1.1$	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.
111.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.
1.11	.	.	.	.	.	.	.	.	.	.	.	.	.	1	.

References

[FS90] P. Fong and B. Srinivasan. Brauer trees in classical groups. *J. Algebra*, 131(1):179–225, 1990.