

$${}^2A_7(q) \bmod l, d = 14$$

[illegible]

	<i>ps</i>	<i>ps</i>	<i>ps</i>	<i>ps</i>	<i>ps</i>	<i>ps</i>
4.	.	.	.	.	.	.
3.1	.	.	.	.	.	.
2.11	.	.	.	.	.	.
1.111	.	.	.	.	.	.
.1111	.	.	.	.	.	.
${}^2A_5 : 2$	.	.	.	.	.	.
${}^2A_5 : 11$	.	.	.	.	.	.
.4	.	.	.	.	.	.
31.	.	.	.	.	.	.
1.3	.	.	.	.	.	.
.31	.	.	.	.	.	.
2.2	.	.	.	.	.	.
22.	.	.	.	.	.	.
21.1	.	.	.	.	.	.
211.	.	.	.	.	.	.
11.2	.	.	.	.	.	.
1.21	1	.	.	.	.	.
.22	.	1	.	.	.	.
.211	.	.	1	.	.	.
11.11	.	.	.	1	.	.
111.1	.	.	.	.	1	.
1111.	.	.	.	.	.	1

References

- [FS90] P. Fong and B. Srinivasan. Brauer trees in classical groups. *J. Algebra*, 131(1):179–225, 1990.